

## CRASH COURSE

# CBSE 12<sup>th</sup> 2021-22

## PHYSICS

※ This is for the **Sample Study Material**. We provides all the study material to purchased user. Please check the course details on [www.misostudy.com](http://www.misostudy.com)

### CBSE 2021-22 CRASH COURSE

CBSE Class 12th PCB crash course provides a preparation strategy & direction, a speedy revision and getting a high score for CBSE Class 12th PCB. It is a focused course for the CBSE aspirant's full preparation through a final mock test with important exam pattern, solving past questions and emphasize on the formulas to crack the CBSE Class 12th exam.

▲ Important problem-solving and revision of all important topics with the last 7 years CBSE Class 12th analysis. ▲ Providing problem-solving tips and tricks for the exam. ▲ 100% CBSE pattern questions with detailed solutions. ▲ Those questions are the focus on chapters with a high weight. ▲ Misconceptions and repeated errors are cleared by the faculties. ▲ The questions of complete syllabus designed by the experienced Misostudy faculty team. ▲ Boosts confidence in students so that they can score well.

### (SUBJECTIVE QUESTIONS)

1. How is force between two charges affected when each charge is doubled and distance between them is also doubled?
2. How much positive and negative charge is there in a cup of water?
3. A metal sphere has a charge of  $-6.5 \mu\text{C}$ . When  $5 \times 10^{13}$  electrons are removed from the sphere, what would be the net charge on it?
4. Define relative permittivity of a medium.
5. Explain the phenomenon of charging by induction?

## Answer & Solutions

1.

As  $F \propto \frac{|q_1||q_2|}{r^2}$   $\therefore$  F becomes  $\frac{(2)(2)}{2^2}$  time = 1 time, i.e., force remains the same.

2.

So in one cup of water (250 ml) there are  $1.338 \times 10^7$  C positive charge and  $-1.338 \times 10^7$  C negative charge. Totally they will be neutral.

3.

So, 8 uC charge transferred from sphere. so, net charge =  $-6.5 + 8 = 1.5$  uC

4.

Relative permittivity  $\epsilon(r)$  is defined as the ratio of absolute permittivity of a medium ( $\epsilon$ ) to the absolute permittivity of free space ( $\epsilon_0$ )

$$\epsilon(r) = \epsilon / \epsilon_0$$

It can also be defined as the ratio of forces of attraction/repulsion between two point charges separated by a certain distance in vacuum to the force of attraction when they are placed in a medium same distance apart.

$$\epsilon(r) = F(o) / F(\text{medium})$$

5.

When charging a conductor by induction, a charged object is brought close to but does not touch the conductor. In the end the conductor has charge of the opposite sign as the charge on the object. One way to carry out the four-step process :

- (i) Bring the charged object close to, but not touching, the conductor. Charge on the conductor shifts in response to the nearby charged object.
- (ii) Connect the conductor to ground. Ground is basically a charge reservoir - anything that can give up or receive charge without noticing the change. Electrons flow from ground to the conductor if the charged object is positive, and the opposite way if the object is negative. The conductor now has a net charge with a sign opposite to the sign on the charged object.
- (iii) Remove the ground connection. The transferred electrons can't get back to where they came from.
- (iv) Remove the charged object. The net charge distributes itself over the surface of the conductor.

**(OBJECTIVE QUESTIONS)**

- Two equal charges placed in air separated by a distance 3 m repel each other with force 0.1 gf. Calculate the magnitude of either of the charges.
  - $9.9 \times 10^{-7} \text{ C}$
  - $4.5 \times 10^{-7} \text{ C}$
  - $12.4 \times 10^{-7} \text{ C}$
  - $15.2 \times 10^{-7} \text{ C}$
- An electron is released with a velocity of  $5 \times 10^6 \text{ ms}^{-1}$  in an electric field of  $10^3 \text{ NC}^{-1}$  which has been applied so as to oppose its motion. The distance travelled by electron before coming to
  - $7.11 \times 10^{-2} \text{ m}$
  - $3.35 \times 10^{-2} \text{ m}$
  - $9.2 \times 10^{-2} \text{ m}$
  - $11.4 \times 10^{-2} \text{ m}$
- Three charges, each equal to  $q$ , are placed at the three corners of a square of side  $a$ . Find the electric field at the fourth corner of the square
  - $\frac{1}{4\pi\epsilon_0} \frac{q}{2a^2}$
  - $\frac{1}{4\pi\epsilon_0} \frac{q}{2a^2} (2\sqrt{2} + 1)$
  - $\frac{1}{4\pi\epsilon_0} \frac{q}{3a^2} (\sqrt{2} + 1)$
  - $\frac{1}{4\pi\epsilon_0} \frac{q}{3a^2}$
- If the number of electric lines of force emerging out of a closed surface are  $10^3$ , calculate the charge enclosed by the surface
 

(a) $4.42 \times 10^{-9} \text{ C}$	(c) $17.7 \times 10^{-9} \text{ C}$
(b) $8.85 \times 10^{-9} \text{ C}$	(d) $12.5 \times 10^{-9} \text{ C}$
- A uniform electric field  $\vec{E} = 5 \times 10^3 \text{ NC}^{-1}$  exist in space. The flux of this field through a square of 10 cm on a side, and whose plane is parallel to the YZ plane is
  - $25 \text{ Nm}^2 \text{ C}^{-1}$
  - $50 \text{ Nm}^2 \text{ C}^{-1}$
  - $75 \text{ Nm}^2 \text{ C}^{-1}$
  - $100 \text{ Nm}^2 \text{ C}^{-1}$

# Answer & Solutions

1. (a)

For  $q_1 = q_2 = q$  (say),

$$F = \frac{1}{4\pi\epsilon_0} \cdot \frac{q \times q}{r^2}$$

$$\text{or } q = \sqrt{4\pi\epsilon_0 F \times r}$$

Setting  $F = 0.1 \text{ gf} = 0.1 \times 10^{-3} \text{ kgf} = 0.1 \times 10^{-3} \times 9.8 \text{ N}$ ,

$$r = 3 \text{ m and } \epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2\text{N}^{-1} \text{ m}^{-1}$$

We get

$$q = 9.9 \times 10^{-7} \text{ C}$$

2. (a)

Since electric field is applied so as to oppose the motion of electron,

$$a = -\frac{eE}{m} = -\frac{1.6 \times 10^{-19} \times 10^3}{9.1 \times 10^{-31}}$$

$$= -1.758 \times 10^{14} \text{ m s}^{-2} \text{ (retardation)}$$

Now,  $u = 5 \times 10^6 \text{ m s}^{-1}$ ,  $v = 0$ 

Using the relation :

$$v^2 - u^2 = 2aS, \text{ we get}$$

$$S = 7.11 \times 10^{-2} \text{ m}$$


3. (b)

Here,  $AB = BC = CD = AD = a$ 

$$\therefore BD = \sqrt{a^2 + a^2} = \sqrt{2}a$$

$$\text{Now, } E_A = E_C = \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{a^2}$$

$$\text{and } E_B = \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{(\sqrt{2}a)^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{2a^2}$$



Since  $E_A$  and  $E_C$  are equal, their resultant will be equally inclined to them i.e it will along  $BD$ .  
If is resultant of  $E_A$  and  $E_C$  them

$$E' = \sqrt{E_A^2 + E_c^2} = \sqrt{E_A^2 + E_A^2} = \sqrt{2} E_A$$

$$= \frac{1}{4\pi\epsilon_0} \cdot \frac{\sqrt{2}q}{a^2}$$

Hence, the resultant of electric fields due to the three charges,

$$E = E + E_B = \frac{1}{4\pi\epsilon_0} \cdot \frac{\sqrt{2}q}{a^2} + \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{2a^2}$$

$$= \frac{1}{4\pi\epsilon_0} \cdot \frac{q}{2a^2} (\sqrt{2} + 1)$$

4. (b)

$$q = \epsilon_0 \Phi$$

$$= 8.854 \times 10^{-12} \times 10^3 = 8.854 \times 10^{-9} \text{ C}$$

5. (b)

$$\text{Flue } \Phi = \vec{E} \cdot \vec{A}$$

$$= (5 \times 10^3 \hat{i}) (10 \times 10 \times 10^{-4}) \hat{i}$$

$$= 50 \text{ Nm}^2 \text{ C}^{-1}$$

miso  
study

# CHEMISTRY

## CBSE 2021-22 CRASH COURSE

CBSE Class 12th PCB crash course provides a preparation strategy & direction, a speedy revision and getting a high score for CBSE Class 12<sup>th</sup> PCB. It is a focused course for the CBSE aspirant's full preparation through a final mock test with important exam pattern, solving past questions and emphasize on the formulas to crack the CBSE Class 12th exam.

▲ Important problem-solving and revision of all important topics with the last 7 years CBSE Class 12th analysis. ▲ Providing problem-solving tips and tricks for the exam. ▲ 100% CBSE pattern questions with detailed solutions. ▲ Those questions are the focus on chapters with a high weight. ▲ Misconceptions and repeated errors are cleared by the faculties. ▲ The questions of complete syllabus designed by the experienced Misostudy faculty team. ▲ Boosts confidence in students so that they can score well.

## (SUBJECTIVE QUESTIONS)

(Q. 1 One Marks)

1.

- (i) “Crystalline solids are anisotropic in nature”. What does this statement mean.
- (ii) Classify the following into different categories of crystalline solids :-  
Urea , ammonia, tin, graphite, silicon, potassium sulphate, water, argon

(Q. 2 Two Marks)

2. Explain the following terms with suitable examples :

- (i) Schottky defect
- (ii) Frenkel defect.

(Q. 3 Three Marks)

3.

- (i) What do you understand by colligative properties ?
- (ii) State Raoult's law for a binary solution containing volatile components.

(Q. 4 Four Marks)

4. Explain the following with suitable examples:

- (i) Ferrimagnetism
- (ii) Paramagnetism
- (iii) Ferromagnetism
- (iv) F-centre.

(Q. 5 Four Marks)

5.

- (i) Define the following terms :
- Isotonic solutions and hypertonic solutions
  - Van't Hoff factor
  - Plasmolysis
- (ii) What is meant by 'reverse osmosis'?



## Answer & Solutions


- 1.
- (i) Anisotropic means that properties of crystalline solids like refractive index thermal conductivity electrical conductivity etc. when calculated from different directions comes out to be different. All Crystalline solids are not Anisotropic. Some Crystalline solids are Anisotropic because despite showing periodicity they are not exactly the same in all directions. It all depends on the symmetry of the unit cell of the crystal. They have their atoms arranged in different manner and in three different plane (X,Y,Z).

(ii)

Potassium sulphate	Ionic solid
Benzene	Molecular solid (non-polar)
Urea	Polar molecular solid
Ammonia	Polar molecular solid
Water	Hydrogen bonded molecular solid
Zinc sulphide	Ionic solid
Graphite	Covalent or network solid
Rubidium	Metallic solid
Argon	Non-polar molecular solid
Silicon carbide	Covalent or network solid

2.


- (i) It is basically a vacancy defect in ionic solids. In order to maintain electrical neutrality, the number of missing cations and anions are equal. Like simple vacancy defect, Schottky defect also decreases the density of the substance. Number of such defects in ionic solids is quite significant.



Schottky defects

- (ii) This defect is shown by ionic solids. In this defect the smaller ions are dislocated from its normal site to an interstitial site. It generates a vacancy defect at its original site and an interstitial defect at its new location. Frenkel defect is also called dislocation defect. In this type of defect, density remains same. Frenkel defect is shown by ionic substance in which there is a large difference in the size of ions, for example, ZnS, AgCl, AgBr and AgI shows this effect due to small size of  $Zn^{2+}$  and  $Ag^+$  ions.






Frenkel defects


- 3.
- (i) that depend on the ratio of the number of solute particles to the number of solvent molecules in a solution, and not on the nature of the chemical species present. ... The word colligative is derived from the Latin colligatus meaning bound together
  - (ii) containing a non-volatile solute is directly related to the mole fraction of solvent (i.e. volatile) in the solution. Relative going-down of vapour pressure is equal to mole fraction of non volatile and non-electrolytic solute.

4.


- (i) Schottky defect is basically a vacancy defect shown by ionic solids. In this defect, an equal number of cations and anions are missing to maintain electrical neutrality. It decreases the density of a substance. Significant number of Schottky defects is present in ionic solids. For example, in NaCl, there are approximately  $10^6$  Schottky pairs per  $\text{cm}^3$  at room temperature. Ionic substances containing similar-sized cations and anions show this type of defect. For example: NaCl, KCl, CsCl, AgBr, etc.




- (ii) Ionic solids containing large differences in the sizes of ions show this type of defect. When the smaller ion (usually cation) is dislocated from its normal site to an interstitial site, Frenkel defect is created. It creates a vacancy defect as well as an interstitial defect. Frenkel defect is also known as dislocation defect. Ionic solids such as AgCl, AgBr, AgI and ZnS show this type of defect.



- (iii) Interstitial defect is shown by non-ionic solids. This type of defect is created when some constituent particles (atoms or molecules) occupy an interstitial site of the crystal. The density of a substance increases because of this defect.



- (iv) When the anionic sites of a crystal are occupied by unpaired electrons, the ionic sites are called F-centres. These unpaired electrons impart colour to the crystals. For example, when crystals of NaCl are heated in an atmosphere of sodium vapour, the sodium atoms are deposited on the surface of the crystal. The Cl ions diffuse from the crystal to its surface and combine with Na atoms, forming NaCl. During this process, the Na atoms on the surface of the crystal lose electrons. These released electrons diffuse into the crystal and occupy the vacant anionic sites, creating F-centres.







5.

(i)

- (a) Mole fraction of a component is the ratio of number of moles of the component to the total number of moles of all the components.
- (b) Van't Hoff factor is the ratio of normal molar mass to the abnormal molar mass. Van't Hoff factor is the ratio of observed value of colligative property to calculated value of colligative property assuming no association or dissociation.

**(OBJECTIVE QUESTIONS)**

1. Which of the following arrangements shows schematic alignment of magnetic moments of antiferromagnetic substances ?

- (a) 
- (b) 
- (c) 
- (d) 

2. Which of the following point defects are shown by AgBr(s) crystals ?

- (A) Schottky defect (B) Frenkel defect  
(C) Metal excess defect  
(D) Metal deficiency defect

- (a) (A) and (B) (c) (A) and (C)  
(b) (C) and (D) (d) (B) and (D)

3. To get an *n*-type semiconductor from silicon, it should be doped with a substance having following valance electrons.

- (a) 2 (c) 3  
(b) 1 (d) 5

4. A compound formed by elements X and Y crystallizes in a cubic structure in which atoms X are at the corners of the cube and atoms Y are at the face centres. The formula of the compound is :

- (a)  $X_3Y$  (c)  $XY_2$   
(b)  $XY$  (d)  $XY_3$

5. AB crystallizes in body centred cubic lattice with edge length 'a' equal to 387 pm. The distance between two oppositely charged ions in the lattice is :

- (a) 300 pm (c) 250 pm  
(b) 355 pm (d) 200 pm

## Answer & Solutions

Q1. (d)

The substances which are expected to possess paramagnetism or ferromagnetism on the basis of magnetic moments of domains but actually they possess zero net magnetic moment are called anti-ferromagnetic substances.

They possess equal no. of domains in opposite directions.

Q2. (a)

Schottky defect is shown by the ionic compounds having comparative size of cation & Anion. Frenkel defect occurs when the cationic size is smaller.

Despite of both the opposite facts, the radius ratio of AgBr being intermediate, makes it a compound showing both defects.

Majorly AgBr shows frenkel defect.

Q3. (d)

Si has 4 valence electrons when it is doped with a compound having 5 valence electrons, it forms 4 covalent bonds with the doped atom, and the 5<sup>th</sup> electron is delocalised, increasing the conductivity of silicon and making it negatively charged. This is how n-type semiconductor is formed.

Q4. (d)

corners  $\rightarrow$  8

$$\therefore \frac{1}{8} \times 8 = 1 \text{ atoms (X)}$$

face centres = 6

$$\frac{1}{2} \times 6 = 3 \text{ atoms (Y)}$$

$XY_3$  is the formula

Q5. (b)

In BCC structure, the cations occupy the corners, whereas the anions occupy the centre of the cube.

$\therefore$  the diagonal of cube  $\sqrt{3} \times a$

$$\text{Distance between 2 oppositely charged ions} = \frac{\sqrt{3}}{2} a = \frac{\sqrt{3}}{2} \times 387 = 335.15 \text{ pm}$$

# BIOLOGY

## CBSE 2021-22 CRASH COURSE

CBSE Class 12th PCB crash course provides a preparation strategy & direction, a speedy revision and getting a high score for CBSE Class 12<sup>th</sup> PCB. It is a focused course for the CBSE aspirant's full preparation through a final mock test with important exam pattern, solving past questions and emphasize on the formulas to crack the CBSE Class 12th exam.

▲ Important problem-solving and revision of all important topics with the last 7 years CBSE Class 12th analysis. ▲ Providing problem-solving tips and tricks for the exam. ▲ 100% CBSE pattern questions with detailed solutions. ▲ Those questions are the focus on chapters with a high weight. ▲ Misconceptions and repeated errors are cleared by the faculties. ▲ The questions of complete syllabus designed by the experienced Misostudy faculty team. ▲ Boosts confidence in students so that they can score well.

## (SUBJECTIVE QUESTIONS)

(Q.1 One Marks)

1. Offsprings produced by asexual reproduction are called clones. Justify giving two reasons.

(Q.2 Three Marks)

2.

- (i) List the three states the annual and biennial angiosperms have to pass through during their life cycle.
- (ii) List and describe any two vegetative propagules in flowering plants.

(Q.3 Three Marks)

3. Draw a diagram of a mature embryo sac of an angiosperm and label the following parts in it.

- (i) Filiform apparatus
- (ii) Synergids
- (iii) Central cells
- (iv) Egg cell
- (v) Polar nuclei
- (vi) Antipodals

(Q.4 Three Marks)

4.

- (i) Name the organic material of the pollen grain is made up of. How is this material advantageous to pollen grain?
- (ii) Still it is observed that it does not form a continuous layer around the pollen grain Give reason.
- (iii) How are 'pollen banks' useful?

(Q.5 Five Marks)

5.

- (i) Describe the sequence of the process of microsporogenesis in angiosperms.
- (ii) Draw a labelled diagram of a 2-celled final structure formed.



## Answer & Solutions

1.

They are called as the clones as they are very much similar to their parents that we can see in their morphology and genetics. All of the individual that reproduce by asexual reproduction are same to their parents as their is no crossing over during the fertilization which ensures the new combinations.

2.

(i) Vegetative propagules are the parts/units of a plant which can be used for vegetative propagation, e.g. roots, stems, leaves, etc.

Root propagules include the production of a new plant via roots, e.g. fleshy roots in case of sweet potato, tapioca and Dahlia.

(ii) Vegetative propagation through leaves, many plant leaves have adventitious buds which help in the development of a new plant, e.g. Begonia, Bryophyllum, etc.

3.




Fig. A Mature embryo sac of an angiosperm

4.

(i) The hard outer layer called exine is made up of sporopollenin which is one of the most resistant organic material. It can withstand high temperature, strong acids and alkalis. It cannot be degraded by any of the known enzymes. Hence, it acts as a shield and protects the pollen grain from getting damaged.

(ii) Exine does not form a continuous layer around the pollen grain. Pollen grain exine has prominent aperture called germ pore where sporopollenin is absent. Germ pores serve as an outlet for the formation of pollen tube.

(iii) Pollen grains at a large can be stored for years in liquid nitrogen ( $-196^{\circ}\text{C}$ ). So, after this treatment they are stored in pollen banks. Such conserved pollen grains can be used in plant breeding programs

5.

(i) Microsporogenesis – It is a process in which microspores are formed from a pollen mother cell through meiosis.


When anther develops the cells of the sporogenous tissue undergo meiotic divisions to form microspore tetrads.

Each sporogenous tissue gives rise to a microspore tetrad and is known as pollen or microspore mother cell (PMC).

Each microspore mother cell divides meiotically and forms 4 haploid cells or tetrad. These each microspore divides into two unequal cells - large vegetative cell and smaller generative cell.

At this 2 - celled stage the pollen grains are shed. Sometimes the generative cell divides mitotically to give rise to two haploid male gametes that are shed at 3 - celled stage.

(ii)



The above diagram shows two celled final structure formed during microsporogenesis in angiosperms.



## (OBJECTIVE QUESTIONS)

1. Stock and scion are used in

- (a) cutting
- (b) grafting
- (c) layering
- (d) micropropagation

2. In ginger, vegetative propagation occurs through.

- (a) rhizome
- (b) offsets
- (c) bulbils
- (d) runners

3. Which of the following pairs is not correctly match?

Mode of reproduction	Example
(a) Offset	— Water hyacinth
(b) Rhizome	— Banana
(c) Binary	— Sargassum
(d) Conidia	— Penicillium

4. Which of the following processes ensures the continuity of life on earth?

- (a) Reproduction
- (b) Respiration
- (c) Digestion
- (d) Growth and development

5. Budding is found in

- 1. Sycon
- 2. Hydra
- 3. Fasciola
- 4. Obelia

## Answer & Solutions

1. (b)

Stock and scion are used in grafting. Grafting is a horticulture technique wherein tissues from one plant are inserted into those of another, so that the two sets of vascular tissues join together.

This technique is most commonly used in asexual propagation of commercially grown plants. In this technique, one plant is selected for its roots and is called the **stock** or **root stock**.

The other plant is selected for its stems, leaves, flowers or fruits and is called the **Scion**. The scion and stock contains the desired genes to be duplicated in future production by the stock and scion plant.

2. (a)

In ginger, vegetative propagation occurs through rhizome. Rhizomes are stems which grow horizontally under the ground.

In ginger, the underground stems are swollen with food reserves. The terminal bud grows upward to produce the flowering shoot and the lateral buds grow out to form new plant.

3. (c)

The plant body of Sargassum is a diploid sporophyte. It does not multiply asexually by means of spores instead, it the only known method of vegetative means, i.e. fragmentation, which is the only known method of vegetative reproduction in the free-floating species of Sargassum.

4. (a)

Reproduction is the process of formation of new individuals of a species from the pre-existing one. It is meant for perpetuation of a species because the older individuals of each species undergo senescence and die.

5. (b)

Hydra reproduces asexually by exogenous budding, a type of vegetative propagation and sexually by formation of gametes. Hydra reproduces by budding, when plenty of food is available.