

CLASS 10

CBSE

PHYSICS

- ▶ Specially designed eBook for complete CBSE syllabus
- ▶ CBSE preparation strategy & direction, speedy revision
- ▶ Chapter-wise important Problem-Solving
- ▶ Theory, Formulas & Diagrams to crack CBSE
- ▶ Updated New Pattern Objective questions
- ▶ Detailed high quality video lectures by experienced faculties
- ▶ CBSE full preparation for Class IX & X



misostudy
connect smart learning


JEE·NEET·AIIMS·CBSE·FOUNDATION
www.misostudy.com

☎ 8929803804 (MON-FRI: 9am-6pm) support@misostudy.com

MISO STUDY INDIA PVT. LTD.
2ND FLOOR 65-A, OMPRO TOWER, KALU SARAI, NEW DELHI, DELHI 110016

Class 10 | Physic

01 Electricity



Misostudy.com



Scientific eBook

This is a sample eBook. The eBook provides chapter-wise theories and examples, formulas, diagrams, and Exam pattern Problem-Solving of the full syllabus.



Complete video lectures

The theory and Problem-Solving in this eBook provide specially designed the video lectures by experienced faculties. Get Free trial and preview on Misostudy.com.

Electricity is an important source of energy

01. Types of Electric Charges

There are two types of electric charges : *positive* charge and *negative* charge. Important property of electric charges.

- (i) Opposite charges (or Unlike charges) attract each other.
- (ii) Similar charges (or Like charges) repel each other.

The SI unit of electric charge is coulomb which is denoted by the letter C.

A Proton possesses a positive charge of 1.6×10^{-19} C whereas an electron possesses a negative charge of 1.6×10^{-19} C.

The SI unit of electric charge 'coulomb' (C) is equivalent to the charge contained in 6.25×10^{18} electrons.

02. Electric Potential

The electric potential (or potential) at a point in an electric field is defined as the work done in moving a unit positive charge from infinity to that point.

03. Electric Potential

The potential difference between two points in an electric circuit is defined as the amount of work done in moving a unit charge from one point to the other point.

$$\text{Potential difference} = \frac{\text{Work done}}{\text{Quantity of charge moved}}$$

04. Electric Current

It is the potential difference between the ends of the wire which makes the electric charges (or current) to flow in the wire.

The electric current is a flow of electric charges (called electrons) in a conductor such as a metal wire.

$$\text{Current, } I = \frac{Q}{t}$$

The SI unit of electric current is ampere.


When 1 coulomb of charge flows through any cross-section of a conductor in 1 second, the electric current flowing through it is said to be 1 ampere.

Current is measured by an instrument called ammeter. Ammeter is always connected in *series* with the circuit in which the current is to be measured.

An ammeter should have very low resistance.

05. How the Current Flows in a Wire

Electric current is a flow of electrons in a metal wire (or conductor).



6. OHM'S Law

At constant temperature, the current flowing through a conductor is directly proportional to the potential difference across its ends. The ratio of potential difference applied between the ends of a conductor and the current flowing through it is a constant quantity called resistance.

We have just seen that : $\frac{V}{I} = R$

or $V = I \times R$

or $\frac{V}{R} = I$

So, Current, $I = \frac{V}{R}$

- (i) the current is directly proportional to potential difference, and
- (ii) the current is inversely proportional to resistance.

If the potential difference across the ends of a conductor is doubled, the current flowing through it also gets doubled, and if the potential difference is halved, the current also gets halved.

07. Factors Affecting the Resistance of a Conductor

The electrical resistance of a conductor (or a wire) depends on the following factors :

- (i) Length of the conductor,
- (ii) Area of cross-section of the conductor (or thickness of the conductor),
- (iii) Nature of the material of the conductor, and
- (iv) Temperature of the conductor.

We will now describe how the resistance depends on these factors.

08. Resistivity

It has been found by experiments that :

- (i) The resistance of a given conductor is directly proportional to its length, That is : $R \propto l$

- (ii) The resistance of a given conductor is inversely proportional to its area of cross-section.
That is :

$$R \propto \frac{1}{A}$$

By combining the relations both equation, we get :

$$R \propto \frac{l}{A}$$

or

$$R = \frac{\rho \times l}{A}$$

09. Combination of Resistances (Or Resistors)

The resistances can be combined in two ways : (i) in series, and (ii) in parallel.

When two (or more) resistances are connected end to end consecutively, they are said to be connected in series.

When two (or more) resistances are connected between the same two points, they are said to be connected in parallel.

10. Resistances (Or Resistors) in Series

The combined resistance of any number of resistances connected in series is equal to the sum of the individual resistances. If a number of resistances R_1, R_2, R_3, \dots etc., are connected in series, then their combined resistance R is given by : $R = R_1 + R_2 + R_3 + \dots$

11. Resistances (Or Resistors) in Parallel

The reciprocal of the combined resistance of a number of resistances connected in parallel is equal to the sum of the reciprocals of all the individual resistances. If a number of resistances R_1, R_2, R_3, \dots etc., are connected in parallel, then their combined resistance R is given by the formula :

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$$

When a number of resistances are connected in parallel then their combined resistance is less than the smallest individual resistance.

12. Electric Power

Electric power is the electrical work done per unit time.

$$\text{Power} = \frac{\text{Work done}}{\text{Time taken}}$$

$$P = \frac{W}{t}$$

13. Unit of Power

The SI unit of electric power is watt. The power of 1 watt is a rate of working of 1 joule per second.

$$1 \text{ watt} = \frac{1 \text{ joule}}{1 \text{ second}}$$

Electric power is the rate at which electrical energy is consumed. When an electrical appliance consumes electrical energy at the rate of 1 joule per second, its power is said to be 1 watt. The rate at which electrical work is done or the rate at which electrical energy is consumed, is called electric power.

14. Heating Effect of Current

When an electric current is passed through a high resistance wire, like nichrome wire, the resistance wire becomes very hot and produces heat. This is called the heating effect of current.

15. Applications of the Heating Effect of Current

- (i) The heating effect of current is utilised in the working of electrical heating appliances such as electric iron, electric kettle, electric toaster, electric oven, room heaters, water heaters.
- (ii) The heating effect of electric current is utilised in electric bulbs (electric lamps) for producing light.
Tungsten metal is used for making the filaments of electric bulbs because it has a very high melting point.
The electric bulb is filled with a chemically unreactive gas like argon or nitrogen (or a mixture of both).
- (iii) The heating effect of electric current is utilised in electric fuse for protecting household wiring and electrical appliances.

CBSE Pattern Exercise (1)

(Q 1 to 2) Very Short Type

1. Give the law of combination of resistances in series.
2. Two resistances X and Y are connected turn by turn:
 - (i) in parallel
 - (ii) in series.In which case the resultant resistance will be less than either of the individual resistances?

(Q 3 to 5) Short Answer Type

3. A battery of 9 V is connected in series with resistors of 0.2 Ω , 0.3 Ω , 0.4 Ω , 0.5 Ω and 12 Ω . How much current would flow through the 12 Ω resistor?
4. Two resistors, with resistances 5 Ω and 10 Ω respectively are to be connected to a battery of emf 6 V so as to obtain :
 - (i) minimum current flowing
 - (ii) maximum current flowing
5. A 4 Ω coil and a 2 Ω coil are connected in parallel. What is their combined resistance? A total current of 3 A passes through the coils. What current passes through the 2 Ω coil?

(Q 6 to 7) Multiple Choice

6. If two resistor of 25 Ω and 15 Ω are joined together in series and then placed in parallel with a 40 Ω resistor, the effective resistance of the combination is :
 - (a) 0.1 Ω
 - (b) 10 Ω
 - (c) 20 Ω
 - (d) 40 Ω
7. A wire of resistance R_1 is cut into five equal pieces. These five pieces of wire are then connected in parallel. If the resultant resistance of this combination be R_2 , then the ratio $\frac{R_1}{R_2}$ is :
 - (a) $\frac{1}{25}$
 - (b) $\frac{1}{5}$
 - (c) 5
 - (d) 25

(Q 8 to 10) High Order Thinking Skills

8. Two resistances when connected in parallel give resultant value of 2 ohm; when connected in series the value becomes 9 ohm. Calculate the value of each resistance.
9. What will be the current drawn by an electric bulb of 40 W when it is connected to a source of 220 V?
10. An electric bulb is rated 220 V and 100 W. When it is operated on 110 V, the power consumed will be :





Answer & Solution

Q1

The combined resistance of any number of resistances connected in series is equal to the sum of the individual resistances. If a number of resistances R_1, R_2, R_3, \dots etc., are connected in series, then their combined resistance R is given by : $R = R_1 + R_2 + R_3 + \dots$

Q2

When they are in parallel combination.

Q3

The current flow is .67 A.

Q4

For minimum current flow in series – For maximum current flow in parallel.
.4 A, 1.8 A.

Q5

$\frac{4}{3} \Omega$ and 2 A.

Q6

20 Ω

Q7

(d) Hint. Resistance of one piece of wire will be $\frac{R_1}{5}$. And

$$\frac{1}{R_2} = \frac{5}{R_1} + \frac{5}{R_1} + \frac{5}{R_1} + \frac{5}{R_1} + \frac{5}{R_1} = \frac{25}{R_1}$$

Q8

3 Ω and 6 Ω .

Q9

In this case we have been given power P and voltage V , so the formula to be used for calculating the current will be :

$$P = V \times I$$

Here, Power, $P = 40$ watts

Voltage, $V = 220$ volts

And, Current, $I = ?$ (To be calculated)

Now, putting these values in the above formula, we get :

$$40 = 220 \times I$$

$$I = \frac{40}{220}$$

$$= \frac{2}{11}$$

Thus, Current, $I = 0.18$ ampere

Q10

In the first case :

Power, $P = 100$ W

Potential difference, $V = 220$ V

And, Resistance, $R = ?$ (To be calculated)

Now, $P = \frac{V^2}{R}$

So, $100 = \frac{(220)^2}{R}$

And $R = \frac{220 \times 220}{100} = 484 \Omega$

This resistance of 484Ω of the bulb will remain unchanged. *In the second case :*

Power, $P = ?$ (To be calculated)

Potential difference, $V = 110$ V

And, Resistance, $R = 484 \Omega$ (Calculated above)

Now, $P = \frac{V^2}{R}$

$$P = \frac{(110)^2}{484} = \frac{110 \times 110}{484} = 25 \text{ W}$$

CLASS 10

CBSE

CHEMISTRY

- ▶ Specially designed eBook for complete CBSE syllabus
- ▶ CBSE preparation strategy & direction, speedy revision
- ▶ Chapter-wise important Problem-Solving
- ▶ Theory, Formulas & Diagrams to crack CBSE
- ▶ Updated New Pattern Objective questions
- ▶ Detailed high quality video lectures by experienced faculties
- ▶ CBSE full preparation for Class IX & X



misostudy
connect smart learning


JEE·NEET·AIIMS·CBSE·FOUNDATION
www.misostudy.com

☎ 8929803804 (MON-FRI: 9am-6pm) support@misostudy.com

MISO STUDY INDIA PVT. LTD.
2ND FLOOR 65-A, OMPRO TOWER, KALU SARAI, NEW DELHI, DELHI 110016

Class 10 | Chemistry

02 Chemical Reactions and Equations



Misostudy.com



Scientific eBook

This is a sample eBook. The eBook provides chapter-wise theories and examples, formulas, diagrams, and Exam pattern Problem-Solving of the full syllabus.




Complete video lectures

The theory and Problem-Solving in this eBook provide specially designed the video lectures by experienced faculties. Get Free trial and preview on Misostudy.com.

Chemical reactions are the processes in which new substances with new properties are formed. during a chemical reaction, atoms of one element do not change into those of another element. only a rearrangement of atoms takes place in a chemical reaction.

- (i) The substances which take part in a chemical reaction are called reactants.
- (ii) The new substances produced as a result of chemical reaction are called products.

When a magnesium ribbon is heated, it burns in air with a dazzling white flame to form a white powder called magnesium oxide. Actually, on heating, magnesium combines with oxygen present in air to form magnesium oxide :




The burning of magnesium in air to form magnesium oxide is an example of a chemical reaction. Souring of milk (when left at room temperature during summer), Formation of curd from milk, Cooking of food Digestion of food in our body, Process of respiration, Fermentation of grapes, Rusting of iron (when left exposed to humid atmosphere), Burning of fuels (like wood, coal, kerosene, petrol and LPG,) burning of candle wax, and Ripening of fruits, are all chemical changes which involve chemical reactions.

01. Characteristics of Chemical Reactions

The important characteristic of chemical reactions are :

- (i) **Evolution of a Gas** : The chemical reaction between zinc and dilute sulphuric acid is characterised by the evolution of hydrogen gas.



- (ii) **Formation of a Precipitate** : When formed the chemical reaction between potassium iodide and lead nitrate is characterised by the formation of a yellow precipitate of lead iodide.




- (iii) **Change in colour** : When citric acid reacts with potassium permanganate solution, then the purple colour of potassium permanganate solution disappears (it becomes colourless). the chemical reaction between citric acid and purple coloured potassium permanganate solution is characterised by a change in colour from purple to colourless.



- (iv) **Change in Temperature** : The chemical reaction between quicklime and water form slaked lime is characterised by a change in temperature (which is rise in temperature) the reaction between quicklime and water to form slaked lime is an exothermic reaction (which means heat producing reaction).
- (v) **Change in State** : When wax is burned (in the form of a wax candle) then water and carbon dioxide are formed


02. Chemical Equations

The method of representing a chemical reaction with the help of symbols and formulae of the substances involved in it is known as a chemical equation.



Balanced and Unbalanced Chemical Equations

A balanced chemical equation has an equal number of atoms of different elements in the reactants and products.



Let us count the number of atoms of all the elements in the reactants and products separately.

	<i>In reactants</i>	<i>In reactants</i>
No. of Zn atoms	1	1
No. of H atoms	2	2
No. of S atoms	1	1
No. of O atoms	4	4

We find that the reactants contain 1 zinc atom and products also contain 1 zinc atom. Reactants contain 2 hydrogen atoms and products also contain 2 hydrogen atoms, Similarly reactants contain 1 sulphur atom and products also contain 1 sulphur atom. and finally, reactants contain 4 oxygen atoms and the products also contain 4 oxygen atoms. Thus, there is an equal number of atoms of different elements in the reactants and products, so the above chemical equation is a balanced equation Since the number of atoms of various elements in reactants and products is equal we can say that a balanced chemical equation has equal masses of various elements in reactants and products.

An unbalanced chemical equation has unequal number of atoms of one or more elements in the reactants and products.

Balancing of Chemical Equations

The number of making the number of different types of atoms equal on both the sides of an equation is called balancing of equation.



	In reactants	In product
No. of H atoms :	2	2
No. of O atoms :	2	1

	In reactants	In product
No. of H atoms :	2	4
No. of O atoms :	2	2




	In reactants	In product
No. of H atoms :	4	4
No. of O atoms :	2	2

This chemical equation contains an equal number of atoms of hydrogen and oxygen on both sides, so this is a balanced equation.

To make equations more informative

- by indicating the “physical states” of the reactants and products.
- by indicating the “heat changes” taking place in the reaction.
- by indicating the “conditions” under which the reaction takes place.

zinc metal reacts with dilute sulphuric acid to form zinc sulphate solution and hydrogen gas. This can be Written as :



02 Chemical Reactions and Equations

Here, zinc metal is a solid so we write Zn(s)

Dilute sulphuric acid is an aqueous solution, so we write H₂SO₄ (aq)


Zinc sulphate is also an aqueous solution, so we write ZnSO₄(aq)

and hydrogen is a gas which is written as H₂(g)

The above equation can now be written as :



Those reactions in which heat is evolved are known as exothermic reactions. For example, when carbon burns in oxygen to form carbon dioxide, a lot of heat is produced in this reaction :



Those reactions in which heat is absorbed are known as endothermic reactions. For example, when nitrogen and oxygen are heated to a very high temperature (of about 3000°C) they combine to form nitrogen monoxide, and a lot of heat is absorbed in this reaction :



03. Types of Chemical Reactions


- (i) **Combination Reactions** : Those reactions in which two or more substances combine to form a single substance, are called combination reactions.

Example magnesium and oxygen combine when heated to form magnesium oxide :




- (ii) **Decomposition Reactions** : Those reactions in which a compound splits up into two or more simpler substances are known as decomposition reactions.


Example : when calcium carbonate is heated it decomposes to give calcium oxide and carbon dioxide :



Example : when electric current is passed through acidified water, it decomposes to give hydrogen gas and oxygen gas. This reaction can be represented as :



We can carry out the electrolysis of water as follows :



(iii) **Displacement Reactions** : Those reactions in which one element take the place of another element in a compound are known as displacement reaction.

Example when a strip of zinc metal is placed in copper sulphate solution, then zinc sulphate solution and copper are obtained :



(iv) **Double displacement reactions** :

Example when silver nitrate solution is added to sodium chloride solution then a white precipitate of silver chloride is formed along with sodium nitrate solution :



(v) **Oxidation and Reduction Reactions** :

Oxidation :

- The addition of oxygen to a substance is called oxidation.
- The removal of hydrogen from a substance is also called oxidation

Reduction :

- The addition of hydrogen to a substance is called reduction.
- The removal of oxygen from a substance is also called reduction.

Oxidising agent :


- The substance which gives oxygen for oxidation is called an oxidising agent.
- The substance which removes hydrogen is also called an oxidising agent.

Reducing agent :

- The substance which gives hydrogen for reduction is called a reducing agent.
- The substance which removes oxygen is called a reducing agent.

The oxidation and reduction reactions are also called redox reactions (in the name 'redox' the term 'red' stands for 'reduction' and 'ox' stands for oxidation). we will now give some examples of oxidation and reduction reactions.

Example when copper oxide is heated with hydrogen, then copper metal and water are formed :



- In this reaction, CuO is changing into Cu. that is oxygen is being removed from copper oxide. Now by definition removal of oxygen from a substance is called reduction, so we can say that copper oxide is being reduced to copper.
- In this reaction H₂ is changing into H₂O that is oxygen is being added to hydrogen. Now by definition addition of oxygen to a substance is called oxidation, so we can say that hydrogen is being oxidised to water.

04. Effects of Oxidation Reactions in Everyday Life

- Corrosion of metals ,and
- Rancidity of food.

Corrosion : Corrosion is the process in which metals are eaten up gradually by the action of air moisture or a chemical (such as an acid) on their surface. corrosion is caused mainly by the oxidation of metals by the oxygen of air. Rusting of iron metal is the most common form of corrosion.

Rancidity : The condition produced by aerial oxidation of fats and oils in foods marked by unpleasant smell and taste is called rancidity. Rancidity spoils the food materials prepared in fats and oils which have been kept for a considerable time and make them unfit for eating.

CBSE Pattern Exercise (1)

(Q 1 to 3) One Mark

1. State whether the following statement is true or false:
A chemical equation can be balanced easily by altering the formula of a reactant or product.
2. What type of reaction is represented by the digestion of food in our body?
3. What is wrong with the following chemical equation?
$$\text{Mg} + \text{O} \rightarrow \text{MgO}$$
Correct and balance it.

(Q 4 to 5) Two Marks

4. Given one example of a decomposition reaction which is carried out:
 - (a) with electricity
 - (b) by applying heat
5. What type of chemical reactions are represented by the following equations?
 - (a) $\text{A} + \text{BC} \rightarrow \text{AC} + \text{B}$
 - (b) $\text{X} \rightarrow \text{Y} + \text{Z}$
 - (c) $\text{PQ} + \text{RS} \rightarrow \text{PS} + \text{RQ}$
 - (d) $\text{A}_2\text{O}_3 + 2\text{B} \rightarrow \text{B}_2\text{O}_3 + 2\text{A}$

(Q 6 to 7) Three Marks

6.
 - (a) What is meant by a chemical reaction? Explain with the help of an example.
 - (b) Give one example each of a chemical reaction characterised by:
 - evolution of gas
 - change in colour
 - formation of a precipitate
 - change in temperature
7.
 - (a) Define a combination reaction.
 - (b) Give one example of a combination reaction which is also exothermic.
 - (c) Give one example of a combination reaction which is also endothermic.

(Q 8 to 10) Five Marks

8. Translate the following statements into chemical equations and then balance the equations:
 - (a) Hydrogen sulphide gas burns in air to give water and sulphur dioxide.
 - (b) Phosphorus burns in oxygen to give phosphorus pentoxide.
 - (c) Carbon disulphide burns in air to give carbon dioxide and sulphur dioxide
 - (d) Aluminium metal replaces iron from ferric oxide, Fe_2O_3 , giving aluminium oxide and iron.

02 Chemical Reactions and Equations

(e) Barium chloride reacts with zinc sulphate to give zinc chloride and barium sulphate.

9.

(a) What is a redox reaction ? Explain with an example.

(b) When a magnesium ribbon burns in air with a dazzling flame and forms a white ash, is magnesium oxidised or reduced ? Why ?

(c) In the reaction represented by the equation :



- name the substance oxidised.
- name the oxidising agent.
- name the substance reduced.
- name the reducing agent.

10. A red-brown metal X forms a salt XSO_4 . When hydrogen sulphide gas is passed through an aqueous solution of XSO_4 , then a black precipitate of XS is formed along with sulphuric acid solution.

(a) What could the salt XSO_4 be ?

(b) What is the colour of salt XSO_4 ?

(c) Name the black precipitate XS.


(d) By using the formula of the salt obtained in (a) above, write an equation of the reaction which takes place when hydrogen sulphide gas is passed through its aqueous solution.

(e) What type of chemical reaction takes place in this case ?



Answer & Solution

1. Carbon dioxide, CO_2
2. Diffusion
3. The heat required to convert a solid into a liquid or vapour, or a liquid into a vapour, without change of temperature. The two types of latent heats are:
 - Latent heat of fusion
 - Latent heat of vaporisation
4. 1 Kg of heat at 100 degree celsius contains more heat because of the following relationship:
 $\text{Water at } 100^\circ\text{C} + \text{Heat} \rightleftharpoons \text{Steam at } 100^\circ\text{C}$
5. The acetone has low boiling point and thus it gets evaporated immediately by absorbing heat from the palm making us feel cold.
6.
 - (a) Temperature, Pressure



- (c) Evaporation can be made faster by following ways:
- Increasing temperature
 - Increasing surface area

- Decreasing humidity in the surrounding area
- Increasing the speed of wind

7.

- The spreading out and mixing of a substance with another, due to the motion of the particles is known as diffusion. Example: The smell of the perfume spreads due to the diffusion of perfume vapours in the air.
- Particles move very quickly in all the directions.
- O₂ and CO₂, both are dissolved in water which is required by marine animals and plants, respectively, for their survival.

8.

- Pollen Grains
- Water
- Brownian Motion
- The fast moving water particles are constantly hitting particles X, causing them to move in a zig-zag manner.
- Robert brown
- The liquid Y is made up of very small particles which are constantly moving.

9.

- (i) Iodine, (ii) Sodium Chloride, (iii) Naphthalene, (iv) Ammonium Chloride
- W, Y, Z
- Y
- Tincture of Iodine
- W

10.

- 273K
- Freezing
- Latent Heat of Freezing
- Melting Latent Heat of Fusion

CLASS 10

CBSE

BIOLOGY

- ▶ Specially designed eBook for complete CBSE syllabus
- ▶ CBSE preparation strategy & direction, speedy revision
- ▶ Chapter-wise important Problem-Solving
- ▶ Theory, Formulas & Diagrams to crack CBSE
- ▶ Updated New Pattern Objective questions
- ▶ Detailed high quality video lectures by experienced faculties
- ▶ CBSE full preparation for Class IX & X



misostudy
connect smart learning

JEE·NEET·AIIMS·CBSE·FOUNDATION
www.misostudy.com

☎ 8929803804 (MON-FRI: 9am-6pm) support@misostudy.com


MISO STUDY INDIA PVT. LTD.
2ND FLOOR 65-A, OMPRO TOWER, KALU SARAI, NEW DELHI, DELHI 110016

Cycle in Mosq

Class 10 | Biology

03 Life Processes

Sporozoites



Misostudy.com



Scientific eBook

This is a sample eBook. The eBook provides chapter-wise theories and examples, formulas, diagrams, and Exam pattern Problem-Solving of the full syllabus.



Complete video lectures

The theory and Problem-Solving in this eBook provide specially designed the video lectures by experienced faculties. Get Free trial and preview on Misostudy.com.

01. Introduction

All the plants and animals (including human beings) are alive or living things. The most important criterion to decide whether something is alive (or not) is the movement. All the living things (which are alive) move by themselves without any external help. The movement in most of the animals are fast and can be observed easily but the movements in plants are usually slow and observed with difficulty. Animals can move from one place to another or they can move their body parts. The plants are fixed in the soil at a place, so they cannot move like animals from place to place. The plants can only move parts of their body such as leaves, flowers, shoots and roots. The plant parts move towards a stimulus such as sunlight, gravity or water, etc. Non-living things (which are not alive) cannot move by themselves. The characteristics of living things are as follows :

- Living things can move by themselves.
- Living things need food, air and water.
- Living things can grow.
- Living things can respond to changes around them. They are sensitive.
- Living things respire (release energy from food).
- Living things excrete (get rid of waste materials from their body).
- Living things can reproduce. They can have young ones.

02. What are Life Processes

The basic functions performed by living organisms to maintain their life on this earth are called life processes. The basic life processes common to all the living organisms are Nutrition and Respiration; Transport and Excretion; Control and coordination (Response to stimuli); Growth; Movement and Reproduction.

03. Energy is Need for The Life Processes

Food is a kind of fuel which provides energy to all the living organisms. The energy required by an organisms comes from the food that the organism eats.

04. Nutrition

Food is an organic substance. The simplest food is glucose. It is also called simple sugar. A more complex food starch. Starch is made from glucose. The general name of substances like glucose (sugar) and starch is 'Carbohydrates are the most common foods for getting energy. Fats and proteins are also foods. (A wider definition of food, however, also includes mineral salts, vitamins and water which are essential for the normal growth and development of an organism). The process of taking in food (consuming food) and utilising it is called nutrition. A nutrient can be defined as a substance which an organism obtains from its surroundings and uses it as source of energy or for the biosynthesis of its body constituents (like tissues and organs). For example carbohydrate and fats are the nutrients. Nutrition is a process of

intake of nutrients (like carbohydrates, fats, proteins, minerals, vitamins and water) by organism as well as the utilisation of these nutrients by the organism.

05. Modes of Nutrition

There are mainly two modes of nutrition :

- (i) **Autotrophic Mode of Nutrition** : Autotrophic nutrition is that mode of nutrition in which an organism make (or synthesizes) its own food from the simple inorganic materials like carbon dioxide and water present in the surroundings (with the help of sunlight energy). The green plants have an autotrophic mode of nutrition. The autotrophic bacteria also obtain their food by the autotrophic mode of nutrition. Those organisms which can make their own food from carbon dioxide and water are called autotrophs. All the green plants are autotrophs. The autotrophic organisms (or autotrophs) contain the green pigment called chlorophyll which is capable of trapping sunlight energy.
- (ii) **Heterotrophic Mode of Nutrition** : Heterotrophic nutrition is that mode of nutrition in which an organism cannot make (or synthesize) its own food from simple inorganic materials like carbon dioxide and water, and depends on other organisms for its food. All the animals have a heterotrophic mode of nutrition. Most bacteria and fungi also have heterotrophic mode of nutrition. Those organisms which cannot make their own food from inorganic substances like carbon dioxide and water, and depend on other organisms for their food are called heterotrophs. All the animals are heterotrophs. The non-green plants (like yeast) are also heterotrophs.


Types of Heterotrophic Nutrition

A heterotrophic organism (or heterotroph) can obtain its food from other organisms in three ways. So, the heterotrophic mode of nutrition is of three types:


- (a) **Saprotrophic Nutrition (or Saprophytic)** : Saprotrophic nutrition is that nutrition in which an organism its food decaying organic matter of dead plants, dead animals and rotten bread, etc. Saprophytes are the organisms which obtain their food from dead plants (like rotten leaves), dead and decaying rotten bread
- (b) **Parasitic Nutrition** : The parasitic nutrition is that nutrition in which an organism derives its food from the body of another living organism (called its host) without killing it. A parasite is an organism (plant or animal) which feeds on another living organism called its host.
- (c) **Holozoic Nutrition** : The holozoic nutrition is that nutrition in which an organism take the complex organic food materials into its body by the process of ingestion, the ingested food is digested and then absorbed into the body cells of the organism.

06. Nutrition in Plants

The process by which green plants make their own food (like glucose) from carbon dioxide and water by sunlight energy in the presence of chlorophyll, is called photosynthesis



The process of photosynthesis takes place in the green leaves of a plant



The extra glucose is changed into another food called starch. This starch is stored in the leaves of the plant. The green plants convert sunlight energy into chemical energy by making carbohydrates (foods).

The photosynthesis takes place in the following three steps :

- (i) Absorption of sunlight energy by chlorophyll.
- (ii) Conversion of light energy into chemical energy, and splitting of water into hydrogen and oxygen by light energy.
- (iii) Reduction of carbon dioxide by hydrogen to form carbohydrate like glucose by utilising the chemical energy (obtained by the transformation of light energy).

Condition Necessary for Photosynthesis


- Sunlight,
- Chlorophyll,
- Carbon dioxide, and
- Water.

Raw Materials for Photosynthesis

The raw materials for photosynthesis are :

- Carbon dioxide, and
- Water.

The green plants take carbon dioxide from air for photosynthesis. The carbon dioxide gas enters the leaves of the plant through the stomata present on their surface.




How the Plants Obtain Water for Photosynthesis

The water required by the plants for photosynthesis is absorbed by the roots of the plants from the soil through the process of osmosis. The water absorbed by the roots of the plants is transported upward through the xylem vessels to the leaves where it reaches the photosynthetic cells and utilised in photosynthesis. The plants take materials like nitrogen, phosphorus, iron and magnesium, etc., from the soil.

Site of Photosynthesis : Chloroplasts

Photosynthesis occurs in the organelles called chloroplasts present in the photosynthesis cells (or mesophyll cells) of green plants. The site of photosynthesis in a cell of the leaf are chloroplasts.



07. Nutrition in Animals

Animals Obtain Their Food From Plants or Other Animals

Animals obtain their food from plants or other animals (which they eat). We (human beings)

are also animals. We obtain the food like wheat, rice, pulses (*dal*), fruits and vegetables from plants.

All the animals can be divided into three groups on the basis of their food habits (or eating habits). These are :

- Herbivores** : Those animals which eat only plants are called herbivores. Goat, Cow, Buffalo, Sheep, Horse, Deer, Camel, Ass, Ox, Elephant, Monkey
- Carnivores** : Those animals which eat only other animals as food are called carnivores. Those animals which eat only the meat (or flesh) of other animals are called carnivores
- Omnivores** : Those animals which eat both, plants and animals, are called omnivores. Some of the example of omnivores are : Man (Human beings), Dog, Crow, Sparrow, Bear, Mynan, and Ant.


08. Different Steps in The Process of Nutrition in Animals

- **Ingestion** : The process of taking food into body is called ingestion.
- **Digestion** : The process in which the food containing large, insoluble molecules is broken down into small, water soluble molecules (which can be absorbed by the body) is called digestion. Most animals use both, physical and chemical methods for digesting (breaking up) the large food molecules. Physical methods include chewing and grinding the food in mouth and chemical methods include the addition of digestive juices (enzymes) to food by the body itself.
- **Absorption** : The process in which the digested food passes through the intestinal wall into blood stream is called absorption.
- **Assimilation** : The process in which the absorbed food is taken in by body cells used for energy, growth and repair, is called assimilation.
- **Egestion** : The process in which the undigested food is removed from the body is called egestion.

09. Nutrition in Amoeba

Amoeba is a unicellular animal. The mode of nutrition in Amoeba is holozoic. The process of obtaining food by Amoeba is called phagocytosis. The various steps involved in the nutrition of Amoeba are : ingestion, digestion, absorption, assimilation, and egestion.

- **Ingestion** : Amoeba has no mouth or a fixed place for the ingestion of food (intake of food). Amoeba ingests food by using its pseudopodia.



- **Digestion** : In amoeba, food is digested in the food vacuole by digestive enzymes. The enzymes from surrounding cytoplasm enter into the food vacuole and break down the food into small and soluble molecules by chemical reactions
- **Absorption** : The digested food present in the food vacuole of Amoeba is absorbed directly into the cytoplasm of Amoeba cell by diffusion.
- **Assimilation** : A part of the food absorbed in Amoeba cell is used to obtain energy through respiration. The remaining part of absorbed food is used to make the parts of Amoeba cell which lead to the growth of Amoeba.
- **Egestion** : Amoeba has no fixed place (like anus) for removing the undigested part of food. When a considerable amount of undigested food collects inside Amoeba, then its cells membrane suddenly ruptures at any place and the undigested food is thrown out of the body of Amoeba.

10. Nutrition in Human Beings (Human Digestive System)

The human digestive system consists of the alimentary canal and its associated glands. The various organs of the human digestive system in sequence are : Mouth, Oesophagus (or Food pipe), Stomach, Small intestine and Large intestine. The glands which are associated with the human digestive system and form a part of the human digestive system are : Salivary gland, Liver and Pancreas

- **Ingestion** : The human beings have a special organ for the ingestion of food. It is called mouth. The food is put into the mouth with the help of hands.
- **Digestion** : The mouth cavity (or buccal cavity) contains teeth, tongue, and salivary glands. The teeth cut the food into small pieces, chew and grind it. So, the teeth help in physical digestion. The salivary glands in our mouth produce saliva. Our tongue helps in mixing this saliva with food. Saliva is a watery liquid so it wets the food in our mouth. The salivary glands help in chemical digestion by secreting enzymes. The human saliva contains an enzyme called salivary amylase which digests the starch present in food into sugar. Thus, the digestion of starch (carbohydrate) begins in the mouth itself. The slightly digested food in the mouth is swallowed by the tongue and goes down the food pipe called oesophagus

The contraction and expansion movement of the walls of food pipe is called peristaltic movement. The peristaltic movement of food pipe (or oesophagus) pushes the slightly digested food into the stomach. The stomach is a J-shaped present on the left side of the abdomen. The food is further digested in the stomach. The stomach wall contains three tubular glands in its walls. The gland present in the walls of the stomach secrete gastric juice. The gastric juice contains three substances : hydrochloric acid, the enzyme pepsin and mucus. In the acidic medium, the enzyme pepsin beings the digestion of proteins present in food to form smaller molecules. Another function of hydrochloric acid is that it kills any bacteria which may enter the stomach with food. The mucus helps to protect the stomach wall from its own secretion of hydrochloric acid. From the stomach, the partially digested food enters the small

intestine. The small intestine in human beings is the site of complete digestion of food (like carbohydrates, proteins and fats).

- (a) The small intestine receives the secretions of two glands : liver and pancreas. Liver secretes bile. Bile is a greenish yellow liquid made in the liver which is normally stored in the gall bladder. Bile is alkaline, and contains salts which help to emulsify or break the fats (or lipids) present in the food. Thus, bile performs two function : (1) makes the acidic food coming from the stomach alkaline so that pancreatic enzymes can act on it, and (2) bile salt break the fats present in the food into small globules making it easy for the enzymes to act and digest them. Pancreas is a large gland which lies parallel to and beneath the stomach Pancreas secretes pancreatic juice which contains digestive enzymes like pancreatic proteins and the enzyme lipase breaks down the emulsified fats.
- (b) The walls of small intestine contain glands which secrete intestinal juice. : The intestinal juice contains a number of enzymes which complete the digestion of complex carbohydrates into glucose proteins into amino acids and fats into fatty acids and glycerol. Glucose, amino acids, fatty acids and glycerol are small, water soluble molecules.
- **Absorption** : After digestion, the molecules of food become so small that they can pass through the walls of the small intestine (which contain blood capillaries) and go into our blood. This is called absorption. The small intestine is the main region for the absorption of digested food. The presence of villi gives the inner walls of the small intestine a very large surface area. And the large surface area of small intestine helps in the rapid absorption of digested food. The digested food which is absorbed through the walls of the small intestine, goes into our blood.
 - **Assimilation** : The blood carries digested and dissolved food all the parts of the body where it becomes assimilated as part of the cells. This assimilated food is used by the body cells for obtaining energy as well as for growth and repair of the body.
 - **Egestion** : A part of the food which we eat cannot be digested by our body. This undigested food cannot be absorbed in the small intestine. So, the undigested food passed from the small intestine into a wider tube called large intestine (it is called large intestine because it is a quite wider tube). The walls of large intestine absorb most of the water from the undigested food (which the help of villi). Due to this, the undigested part of food becomes almost solid. The last part of the large intestine called 'rectum' stores this undigested food for some time. And when we go to the toilet, then this undigested food is passed out (or egested) from our body through anus as faeces or 'stool' The act of expelling the faeces is called egestion or defecation.

11. Dental Caries

The hard, outer covering of a tooth is called enamel Tooth enamel is the hardest material in our body. The formation of small cavities (or holes) in the teeth due to the action of acid – forming bacteria and improper dental care is called dental caries. If the teeth are not cleaned regularly, they become covered with a sticky, yellowish layer of food particles and bacteria cells called 'dental plaque'

12. Respiration

The assimilated food is used mainly for two purposes :

- (i) Assimilated food is used as a fuel to get energy for various life processes, and
- (ii) Assimilated food is used as a material for the growth and repair of the body.

The process of releasing energy from is called respiration. The process of respiration oxygen (of air) into the cells, using it for releasing energy by burning food, and then eliminating the waste products (carbon dioxide and water) from the body.




(a) Breathing and Respiration

The mechanism by which organisms obtain oxygen from the air and release carbon dioxide is called breathing. Respiration is a more complex process. Respiration includes breathing as well as the oxidation of food in the cells of the organism to release energy.

(b) How Energy Released During Respiration is Stored

The energy produced during respiration is stored in the form of ATP molecules in the cells of the body. ADP is a substance called Adenosine Di-Phosphate. The molecules of ADP are present in a cell. ADP has low energy content. ATP is a substance called Adenosine Tri-Phosphate. ATP has high energy content.

- The energy released during respiration is used to make ATP molecules from ADP and inorganic phosphate.



- When the cell needs energy, then ATP can be broken down using water to release energy. Thus :




The energy equivalent to 30.5 kJ/ mole is released in this process. The energy stored in ATP used by the body cells for various purposes like contraction of muscles, conduction of nerve impulses, synthesis of proteins, and many other activities related to the functioning of cells.


An Important Discussion

- Glucose is $\text{C}_6\text{H}_{12}\text{O}_6$. It is a six carbon atom compound. It is the simple food which is oxidised in the cells of organisms during respiration.
- The oxidation of glucose to pyruvic acid (or pyruvate) is called glycolysis. It occurs in the cytoplasm of a cell and not in mitochondria. The oxidation of glucose to pyruvic acid does not require oxygen. One molecule of glucose on glycolysis produces two molecules of pyruvic acid.

- Pyruvic acid is a three carbon atom compound. It is also called pyruvate. The formula of




- The fate of pyruvate formed during respiration depends on whether oxygen is present in the cells or not. If oxygen is present in the cells, then pyruvate is completely oxidised to carbon dioxide and water, and a lot of energy is produced (in the form of ATP). If, however, oxygen is not present in the cells (that is, in the absence of oxygen), pyruvate is converted to either 'ethanol and carbon dioxide' or 'lactic acid'
- Lactic acid is also a three carbon atom compound. It is also called lactate acid or lactate



13. Types of Respiration


We have two types of respiration : aerobic respiration and anaerobic respiration.

- (a) **Aerobic Respiration** : The respiration which uses oxygen is called aerobic respiration. Aerobic respiration produces a considerable amount of energy for use by the organism which gets stored in the ATP molecules.




Please note that during aerobic respiration (shown above), 1 molecule of glucose (food) produces 38 energy – rich ATP molecules. Mitochondria are the sites of aerobic respiration in the cells.

- (b) **Anaerobic Respiration** : The respiration which takes place without oxygen is called anaerobic respiration. In anaerobic respiration, the micro – organisms like yeast break down glucose (food) into ethanol and carbon dioxide, and release energy.



Please note that

- (c) during anaerobic respiration (shown above), 1 molecule of glucose (food) produces only 2 energy- rich ATP molecules. Anaerobic respiration takes place in our muscles during vigorous physical exercise when oxygen gets used up faster in the muscle cells than can be supplied by the blood.



The sudden build up of lactic acid in our muscle during vigorous physical activity can cause muscular 'cramps'. We can get relief from cramps in muscles caused by heavy exercise by taking a hot water bath or a massage.

Differences between Aerobic and Anaerobic Respiration

Aerobic respiration	Anaerobic respiration
<ul style="list-style-type: none"> Aerobic respiration takes place in the presence of oxygen Complete breakdown of food occurs in aerobic respiration The end products in aerobic respiration are carbon dioxide and water. Aerobic respiration produces a considerable amount of energy. 	<ul style="list-style-type: none"> Anaerobic respiration place in the absence of oxygen. Partial breakdown of food occurs in anaerobic respiration. The end products in anaerobic respiration may be ethanol and carbon dioxide (as in animal muscles). Much less energy is produced in anaerobic respiration.

14. Respiration in Plants

The respiration in plants differs from that in animals in three respects :

- All the parts of a plant (like root, stem and leaves) perform respiration individually. On the other hand, an animal performs respiration as a single unit.
- During respiration in plants, there is a little transport of respiratory gases from one part of the plant to the other. On the other hand, respiratory gases are usually transported over long distances inside an animal during respiration.
- The respiration in plants occurs at a slow rate. On the other hand, the respiration in animals occurs at a much faster rate.

Plants get Oxygen by Diffusion

- Respiration in Roots** : Air is present in- between the particles of soil The roots of a plant take the oxygen required for respiration from the air present in- between the soil particles by the process of diffusion. The extensions of the epidermal cells of a root are called root hair. The root hair are in contact with the air in the soil. Oxygen (from air in the soil particles) diffuses into root hairs and reaches all the other cells of the root respiration.
- Respiration in Stems** : The stems of herbaceous plants (or herbs) have stomata. So, the exchange of respiratory gases in the stems of herbaceous plants take place through stomata. In woody stems, the bark (outer covering of stem) has lenticels for gaseous exchange
- Respiration in Leaves** Oxygen from air diffuses into a leaf through stomata and reaches all the cells where it is used in respiration. in leaves occurs during the day time as well as at night. On the other hand, photosynthesis occurs only during the day time
 - During day time, when photosynthesis occurs, oxygen is produced. The leaves use some of the this oxygen for respiration and the rest of oxygen diffuses out into air. Again, during day time, carbon dioxide produced by respiration is all used up in photosynthesis by leaves. The net gas exchange in leaves during day time is : O_2 diffuses out ; CO_2 diffuses in
 - At nigh time when no photosynthesis occurs and hence no oxygen is produced the net gas exchange in leaves at night : O_2 diffuses in ; CO_2 diffuses out.

15. Respiration in Animals

Different animals have different modes of respiration. For example :

- (i) In simple unicellular animals like Amoeba, respiration take place by the simple diffusion of gases through the cell membrane. Most of the animals have, specific organs for respiration
- (ii) The animals like earth worms which live in the soil use their skin to absorb oxygen from air and remove carbon dioxide. So, the respiratory organ in the earthworm is the skin.
- (iii) The aquatic animals like fish, prawns and mussels have gills as the respiratory organs which extract oxygen dissolved in water and take away carbon dioxide from the body.
- (iv) In the insects like grasshopper, cockroach, housefly and a mosquito, the tiny holes called spiracles on their body and the air tubes called tracheae are the respiratory organs
- (v) The respiratory organs of the land animals such as man (humans), birds, lizard, dog, and frog, etc., are the lungs. (Frogs, however, breathe both by lungs and skin).

Respiration in Human

The process by which energy is released from food in our body is called respiration. The function of respiratory system is to breathe in oxygen for respiration (producing energy from food), and to breathe out carbon dioxide produced by respiration. Breathing is the process by which air in oxygen is taken inside the body of an organism and air rich in carbon dioxide is expelled from the body. The taking in of air rich in oxygen into the body during breathing is called *inhalation* giving our (or expelling) the air rich in carbon dioxide is *exhalation* ` A breath means` one inhalation plus one exhalation`

- (a) **reathing in** : When we breathe in (or inhale), then two things happen at the same time :
 - (i) the muscles between the ribs contract causing the rib cage to move upward and outward, and (ii) the diaphragm contracts and move downward The upward and outward movement of rib cage, as well as the downward movement of diaphragm, both increase the space in the chest cavity and make it larger As the chest cavity becomes larger, air is sucked in from outside into the lungs. The lungs get filled up with air and expand
- (b) **Breathing out** : When we breathe our (or exhale), even then two things happen at the same time : (i) the muscles between the ribs relax causing the rip cage to move downward and inward, and (ii) the diaphragm relaxes and move upward. The downward and inward movement of rib cage, as well as the upward movement of diaphragm, both decrease the space in chest cavity and make it smaller as the chest cavity becomes smaller, air is pushed out from the lungs.

Respiration in Plants

- All parts of a plant perform respiration.
- Plants exchange gases by diffusion through the stomata.
- Oxygen from the air diffuses into a leaf and reaches all the cells for respiration.
- Carbon dioxide produced during respiration is released into the air through the stomata.
- In plants, respiration occurs during the day as well as during the night.
- During the day, oxygen produced during photosynthesis is used for respiration and the extra amount of oxygen is given out through the stomata.
- The roots of plants take up oxygen from the air present between the roots and soil particles.
- In stems, the exchange of gases occurs through either the stomata or lenticels.


Respiration in Animals

Different animals have evolved different respiratory organs:



Human Respiratory System

The respiratory system in human beings consists of the nose, pharynx, larynx, trachea, bronchi and lungs.





- Air is taken in through the nostrils.
- Hairs present in the nostrils prevent the entry of dust particles inside the nose.
- Nostrils further continue into the nasal cavity.
- Nostrils humidify the air passing through it.
- There is a bony plate called the palate, which separates the oral cavity from the nasal cavity.
- Nasal cavity opens into the pharynx.
- The pharynx is a muscular chamber acting as a common passage for the windpipe or trachea and the food pipe or oesophagus.
- It is connected to the larynx through a slit-like opening called the glottis.
- The larynx is also called the voice-box or Adam's apple.
- The larynx connects the pharynx to the trachea.
- The trachea shows the presence of cartilaginous rings.
- The cartilaginous rings provide flexibility thus, facilitating continuous air flow.
- The inner wall of the trachea is lined by a mucous membrane consisting of ciliated columnar epithelium.

- The trachea divides into two branches or tubes called bronchi, one of which enters the right lung and the other enters the left lung.
- The bronchi have cartilaginous rings for distention.
- Each bronchus divides into fine secondary bronchi. These bronchi further divide into finer tertiary bronchi. In the lungs, each bronchus finally divides into finer and smaller branches called bronchioles.
- The bronchioles further divide to form smaller terminal bronchioles.
- The bronchioles divide repeatedly to form a cluster of tiny air chambers called air sacs or alveoli.
- Alveoli have thin and moist walls which enable gaseous diffusion with blood capillaries.
- The lungs are a pair of spongy and elastic respiratory organs protected by a bony rib cage.
- The base of the lungs rests on the diaphragm.
- Each lung is covered by two membranes. The inner membrane is called the inner or visceral pleura and the outer membrane is called the outer or parietal pleura.
- The diaphragm is a curved, musculo-fibrous sheath which separates the thoracic cavity from the abdominal cavity.
- The diaphragm plays a major role during respiration.
- The intercostal muscles found between the ribs and the radial muscles of the diaphragm bring about the breathing movements.
- When we breathe in, the ribs are pulled upwards and the diaphragm becomes flat which results in an increase in the volume of lungs.
- When we breathe out, the ribs come back to their normal position, the diaphragm is relaxed, lungs attain their normal size and air is expelled out of the body through the nostrils.

16. Transportation in Plants

- Transportation in plants is the process by which a substance, absorbed or synthesised in one part of the plant, is transported to the other parts of the plant.
- Substances transported by the transport system are water, mineral and food prepared by plants.





(i) **Xylem**

- It conducts water upwards in a plant.
- Xylem also provides strength to the stem and helps the plant to stand upright.
- It is located in the centre of the plant body.
- Xylem mainly consists of tracheids and vessels.

Mechanism of Transport of Water and Minerals

(a)

- Water enters the root hair through osmosis, and mineral ions enter the root cells by active transport.
- Both water and minerals move upward from cell to cell through the cortex of the root by osmosis.
- From the cortex, water and minerals are brought to the xylem.
- The sap which contains water and dissolved minerals move upwards from the root cells to xylem. The upward movement of sap is called the ascent of sap.
- The xylem vessels of the roots are in continuation with the xylem vessels of the stem.

(b)

- Transpiration is the loss of water in the form of water vapour from the aerial parts of a plant.
- It occurs through openings called stomata.
- Water loss through evaporation lowers the concentration of water inside the mesophyll cells.
- Due to this, water enters mesophyll cells from neighbouring xylem vessels through osmosis.
- As water evaporates from the leaves, a suction force is created. This force helps to draw more water up through the stem which causes the roots to absorb more water from the soil.
- Higher the rate of transpiration, greater the rate of absorption of water and solutes from the soil.
- Transpiration also helps in maintaining the temperature of the plant body.

(ii) **Phloem**

- It conducts manufactured food from the leaves to different parts of the plant.
- The food in the phloem can move in the upward and downward directions.
- Phloem mainly consists of sieve tube cells and companion cells.
- Sieve tubes are living cells of the phloem. They contain cytoplasm but no nucleus.
- The end walls of the cells form sieve plates.
- Sieve plates have small pores in them which allow food to pass through the phloem.
- Each sieve tube cell has a companion cell next to it.

Mechanism of Transport of Water and Minerals

- Food synthesised during photosynthesis is loaded into sieve tubes by utilising ATP.
- The presence of food inside the phloem develops the concentration gradient for water. Thus, water enters the phloem by osmosis.
- Osmosis develops high pressure inside the phloem which transports the food from the phloem to plant parts where the concentration of food is less.
- This process is called translocation.
- In spring, the sugar stored in the root or stem tissues is transported to the buds.

17. Blood

Blood is a red coloured liquid which circulates in our body. Blood is red because it contains a red pigment called haemoglobin in its red cells. Blood is a connective tissue. Blood consists of four things : plasma, red blood corpuscles, white blood corpuscles, and platelets. Thus, the main components of blood are :

Blood is a liquid (or fluid matrix) called plasma with red cells, white cells and platelets in it

- Plasma** : The liquid part (or fluid part) of blood is called plasma.
- Red Blood Cells** : Red blood cells carry oxygen from the lungs to all the cells of the body. It is actually the haemoglobin present in the red blood cells which carries oxygen in the body. Red blood cells do not nuclei. Each red blood cell lives for about four months.
- White Blood Cells** : White blood cells fight infection and protect us from diseases This is because white blood cells help to fight against germs and other foreign bodies which cause diseases. White blood cells are called soldiers of the body.
- Platelets** : Platelets help in the coagulation of blood (or clotting of blood) in a cut or wound.

Functions of Blood

- Blood carries oxygen from the lungs to different part of the body.
- Blood carries carbon dioxide from the body cells to the lungs for breathing out.
- Blood carries digested food from the small intestine to all the part of the body.
- Blood carries hormones from the endocrine glands to different organs of the body (where they are needed).
- Blood carries a waste product called urea from the liver to the kidneys for excretion in urine

- Blood protects the body from diseases. This is because white blood cells kill the bacteria and other germs which cause diseases.
- Blood regulates the body temperature. This is because the blood capillaries in our skin help to keep our body temperature constant at about 37°C

Transport in Humans

The main transport system in human beings (or man) is the 'blood circulatory system'. The human blood circulatory system consists of the heart (the organ which pumps and receives the blood) and the blood vessels (or tubes) through which the blood flows in the body.

In blood circulatory system, the blood flows through three types of blood vessels :

- arteries,
- veins, and
- capillaries


In addition to the blood circulatory system for the transport in human beings, there is another system called lymphatic system which also helps in the human body. The liquid which circulates and carries materials in the lymphatic system is called lymph. Thus, in human beings, the various substances are transported through two liquid called 'blood' and 'lymph'

18. Human Circulatory System

The various organs of the circulatory system in humans are : Heart, Arteries, Veins and Capillaries. The heart is roughly triangular in shape. It is made of special muscle called cardiac muscle. The size of our heart is about the same our 'clenched fist'. The heart has four compartments called 'chambers' inside it. The upper two chambers of heart are called atria (Singular atrium), and the lower two chambers of heart are called ventricles. The two atria receive blood from the two main veins. And the two ventricles transport blood to the entire body and the lungs. The left atrium is connected to the left ventricle through a valve V_1 . Similarly, the right atrium is connected to the right ventricle through another valve V_2 . These valves prevent the backflow of blood into atria when the ventricles contract to pump blood out of the heart to the rest of the body. This is because when the ventricles contract, the valves V_1 and V_2 close automatically so that the blood may not go back into atria. The job of heart is to pump blood around our body. All the atria and ventricles of the heart contract and relax (expand) at appropriate times and make the heart behave like a pumps. Since ventricles have to pump blood into various organs with high pressure, they have thicker walls than atria. A sheath of tissue called 'pericardium' protects the muscular heart. The chambers of the heart are separated by a partition called by a partition called septum. The arteries, veins and capillaries are a kind of thin pipes (or tubes) through which blood flows in the body. Arteries are the thick walled blood vessels which carry blood from the heart to all the parts of the body. Arteries are the thick walled blood vessels which carry blood from the heart to all the parts of the body. Arteries have thick walls because blood emerges from the heart under high pressure. Arteries are found in the whole of our body. The main artery (called aorta) is connected to the left ventricle of the heart through a valve V_3 . The main artery carries oxygenated blood from the

left ventricle to all the parts of the body (except the lungs). Another artery called pulmonary artery is connected to the right ventricle of the heart through another valve V_4 . The pulmonary artery carries deoxygenated blood from the right ventricle to the lungs. The capillaries are thin walled and extremely narrow tubes or blood vessels which connect arteries to veins. The exchange of various materials like oxygen, food, carbon dioxide, etc., between the blood and the body cells takes place through capillaries. Veins are the thin walled blood vessels which carry blood which carry blood from all the parts of the body back to the heart.

Veins do not need thick walls because the blood flowing through them is no longer under high pressure. Veins have valves in them which allow the blood in them to flow in only one direction (towards the heart). The valves prevent the backflow of blood in veins. Veins are also found in the whole of our body. The pulmonary vein is connected to the left atrium of the heart. The pulmonary vein carries oxygenated blood from lungs back to the heart. There is also a main vein (called vena cava). The blood from all the parts of the body (except lungs), back to the heart. Please note that the main difference between an artery and a vein is that an artery carries blood from the heart to the body organs whereas a vein carries blood from the body organs back to the heart. The blood carrying oxygen in it is called oxygenated blood. The blood having no oxygen in it is called deoxygenated blood. The heart non-stop all the time.



A circulatory system in which the blood travels twice through the heart in one complete cycle of the body is called double circulation. In the human circulatory system the pathway of blood from the heart to the lungs and back to the heart is called pulmonary circulation; and the pathway of blood from the heart to the rest of the body and back to the heart is called the systemic circulation. The amphibians (like frogs) and reptiles (like lizards) have a three-chambered heart (which consists of two atria and one ventricle). The fish has a two-chambered heart (which consists of one atrium and one ventricle).

Heart Beats : One complete contraction and relaxation of the heart is called a heart beat. The heart usually beats about 70 to 72 times in a minute when we are resting.

Pulse : The expansion of an artery each time the blood is forced into it, is called pulse. The pulse rate of an adult person while resting is 70 to 72 per minute.

Blood Pressure : The pressure at which blood is pumped around the body by the heart is called blood pressure. The blood pressure of a person is always expressed in the form of two values called 'systolic pressure' and diastolic pressure. The maximum pressure at which the blood leaves the heart through the main artery (aorta) during contraction phase, is called the systolic pressure. The minimum pressure in the arteries during the relaxation phase of heart is called the diastolic pressure. The normal blood pressure values are :

- Systolic pressure : 120mm Hg
- Diastolic pressure : 80mm Hg
- This is usually written as 120/80

High blood pressure is called hypertension


How to Measure Blood Pressure

Blood pressure is measured by using an instrument called sphygmomanometer

19. Lymphatic System


Lymph and Lymphatic System

- As the blood flows through capillaries, the water, dissolved substances and a few white blood cells
- pass through the capillary walls into the spaces between the cells, i.e. intercellular spaces. This fluid is called **tissue fluid**.
- White blood cells in the lymph protect the body against diseases.
- The lymphatic system carries excessive tissue fluid back to the blood.




Clotting of Blood (Coagulation)

- When a blood vessel is cut, blood escapes from it. Soon a clot is formed on the wound, and the flow is stopped.
- Blood clotting is a complex process:




20. Excretion

- (i) Excretion is the removal of harmful and unwanted substances, especially nitrogenous wastes, from the body.
- (ii) The human urinary system consists of -




<p>Pair of kidneys</p>	<ul style="list-style-type: none"> • Dark red, bean-shaped, 10 cm long, 6 cm wide. • The right side of the kidney is slightly lower in position due to the presence of the liver.
<p>Pair of ureters</p>	<ul style="list-style-type: none"> • Ureters are tube-like structures which arise from the notch, i.e. the hilum of each kidney. • The ureters connect behind with the urinary bladder. • The ureters carry the urine produced to the urinary bladder
<p>Urinary bladder</p>	<ul style="list-style-type: none"> • Muscular sac-like structure. • It stores urine temporarily. • Its opening is guarded by muscular sphincters. • The sphincters open at the time of micturition (urination).
<p>Urethra</p>	<ul style="list-style-type: none"> • Short muscular tube which expels urine out of the body. • The urethra is long in males and is very short in females. • The opening is guarded by sphincters which open at the time of urination

Urineriferous Tubule



Urineriferous Tubule

- Each kidney is composed of an enormous number of urineriferous tubules.
- They are also known as nephrons, renal tubules or kidney tubules.
- Urineriferous tubules are the structural and functional units of the kidney.



Malpighian Tubule

- Each nephron has a Malpighian body and body of tubules.
- Malpighian body is nothing but a cup-shaped Bowman's capsule. In its up-shaped depression, a tuft of blood capillaries called glomerulus is situated.
- The body of tubules contains proximal convoluted tubule (PCT), loop of Henle and distal convoluted tubule (DCT).
- DCT opens into the collecting duct.

Note :

Approximately 2 million urineriferous tubules are present in both the kidneys.

Each single urineriferous tubule is 4 to 5 cm long.

The great length of the urineriferous tubule provides a large surface area for the reabsorption

of usable substances such as water.

Blood flow through the kidneys per minute = 1 litre

Glomerular filtrate produced in 24 hours = 160 litre

Urine produced from the glomerular filtrate after reabsorption per day = 1.2 litre

Formation of Urine :


The process of urine formation occurs in two major steps.

Ultrafiltration	Reabsorption
<ul style="list-style-type: none"> The efferent arteriole is narrower than the afferent arteriole which develops a hydrostatic pressure on the blood. Thus, the blood flows through the glomerulus with a great pressure. Due to the pressure, the liquid part of the blood filters out from the glomerulus and passes into the Bowman's capsule. The glomerular filtrate consists of water, urea, salts, glucose and other plasma solutes. Blood corpuscles, proteins and other large molecules remain behind in the glomerulus. Therefore, the blood carried away by the efferent arteriole is relatively thick. 	<ul style="list-style-type: none"> The glomerular filtrate entering the renal tubule contains many useful substances. Hence, as the filtrate passes down the tubule, water and other substances required by the body are reabsorbed. Potassium ions and certain substances such as penicillin are passed into the forming urine through the distal convoluted tubule (DCT). The cells of the walls of DCT are involved in bringing potassium ions and other substances back into the renal tubule; hence, this process is known as tubular secretion.

Urine Excretion

- The filtrate left after reabsorption and tubular secretion is called urine.
- The urine passes from the collecting duct into the pelvis of the kidney. From there it is sent to the urinary bladder through the ureters.
- By relaxing the sphincters present at the opening of the urethra, the urine is expelled from the body. This process is known as micturition or urination.

Artificial Kidney



- If one kidney is damaged or removed, then the other kidney alone can fulfil excretory needs.
- However, the failure of both the kidneys allows urea and other wastes to accumulate in the blood.

- Such a patient undergoes dialysis.
- In dialysis, an artificial kidney is used.
- The artificial kidney contains tubes with a semi-permeable lining.
- These tubes are suspended in a tank filled with a dialysing solution.
- This fluid contains water and glucose in concentrations similar to those in blood.
- The patient's blood is led from the radial artery through the tubes of the artificial kidney where excess salts and urea are removed.
- The purified blood is returned through a vein in the same arm.
- The function of dialysis is similar to the function of the kidney, but the only difference is there is no reabsorption during dialysis.

Excretion in Plants

- Plants also produce several waste products during their life processes.
- The major waste products are water, carbon dioxide and oxygen produced during respiration and photosynthesis.
- These wastes are excreted through the stomata and lenticels.
- Plants store some waste products in leaves which fall off.
- Wastes such as gums and resins are stored in the old xylem.

miso
study

CBSE Pattern Exercise (1)

(Q 1 to 3) One Mark

1. Into how many groups can organisms be classified?
2. Name the structures through which amoeba capture food.
3. What is the range of normal systolic and diastolic blood pressure.?

(Q 4 to 6) Two Marks

4. Write one function each of the following components of the transport system in human beings:
 - Blood vessels
 - Blood platelets
 - Lymph
 - Heart.
5. What is the difference between breathing and respiration?
6. Although bile juice has no digestive enzyme, it is still considered to be very important during digestion of food. Give two reasons.

(Q 7 to 8) Three Marks

7. Name the various digestive glands associated with the human digestive system, also give the name of secretion of these glands along with their functions.
8. Why does the medium become acidic in the mouth? What is the ill effect of the acidic medium? How can this be prevented?

(Q 9 to 10) Five Marks

9.
 - (a) Draw the diagram of the human heart and label the following:
 - part which receives deoxygenated blood from the vena cava.
 - part which sends deoxygenated blood to the lung through the pulmonary artery.
 - part which receives oxygenated blood from the lungs.
 - part which sends oxygenated blood to all parts of the body through the aorta.
 - (b) What does the blood consist of?
 - (c) Write two functions of blood.
10. Describe the process of digestion of food in man.



Answer & Solution

Q1.

Two groups :

- Autotrophs and
- Heterotrophs.

Q2.

Pseudopodia.

Q3.

Systolic–120 mm of H.g Diastolic –80 mm of H.g

Q4.

- Blood vessels** : Arteries are the vessels which transport blood from heart to various organs of the body while veins bring blood from various organs of the body back to the heart.
- Blood platelets** : Help in clotting of blood by releasing platelet factor at the site of injury:
- Lymph** : Transports fat and bring back extracellular fluid into circulation.
- Heart** : It pumps oxygenated blood to various organs of the body and deoxygenated blood to lungs for oxygenation.

Q5.

Differences between breathing and respiration:

Breathing	Respiration
<ul style="list-style-type: none"> The mechanism by which an organism obtains oxygen from the environment and release carbon dioxide is termed breathing. 	<ul style="list-style-type: none"> It involves the mechanism of exchange of gases—O_2 and CO_2.
<ul style="list-style-type: none"> Breathing is a physical process. 	<ul style="list-style-type: none"> Oxidation of food occurs in the cells resulting in release of energy which is utilised for carrying out various functions.

Q6

- (i) Bile salts break fats into smaller globules (Emulsification) and thus increase the efficiency of enzyme action.
- (ii) Bile is alkaline in nature. It neutralises the acidic chyme and provides alkaline pH for enzymes secreted by pancreas.

Q7.

The various digestive glands associated with human digestive system are:


<i>Glands</i>	<i>Secretion</i>	<i>Function of secretion</i>
• Salivary glands	Saliva which contains enzyme ptyalin/salivary amylase.	Ptyalin digests starch and converts it to sugar maltose. Mucus present in saliva makes the food slippery so that it can be easily swallowed.
• Gastric glands	HCl and Pepsin	HCl makes the food acidic and pepsin digests protein.
• Liver	Bile juice	Emulsifies fat and bicarbonates present in it turn the acidic food into alkaline.
• Pancreas	Pancreatic juice. It contains enzyme trypsin, lipase and amylase.	Trypsin digest protein. Lipase digest fat and amylase digest starch.
• Intestinal glands	succus entericus	Completes the digestion of starch, protein and fat.

Q8.

The medium in the mouth becomes acidic when bacteria acting on sugars produce acids. The bacterial cells together with food particles stick to the tooth to form dental plaque. As causes tooth decay due to gradual softening of tooth enamel. It can be prevented by brushing the teeth after each meal.

Q9.

(a)



- Right atrium receives deoxygenated blood from vena cava.
 - Right ventricle pumps deoxygenated blood to lungs through pulmonary artery.
 - Left atrium receives oxygenated blood from lungs.
 - Left ventricle sends blood to all parts of body through aorta.
- (b) Blood consists of plasma in which three types of blood cells WBC, RBC and platelets are found. Blood is a type of connective tissue.
- (c)
- Transport of nutrients.
 - Transport of hormones.
 - Transport of excretory products (urea) to kidney for excretion.
 - Transport of gases.

Q10.

Process of digestion starts in mouth

CLASS 10

CBSE

MATHEMATICS

- ▶ Specially designed eBook for complete CBSE syllabus
- ▶ CBSE preparation strategy & direction, speedy revision
- ▶ Chapter-wise important Problem-Solving
- ▶ Theory, Formulas & Diagrams to crack CBSE
- ▶ Updated New Pattern Objective questions
- ▶ Detailed high quality video lectures by experienced faculties
- ▶ CBSE full preparation for Class IX & X



misostudy
connect smart learning

JEE·NEET·AIIMS·CBSE·FOUNDATION
www.misostudy.com


☎ 8929803804 (MON-FRI: 9am-6pm) support@misostudy.com

MISO STUDY INDIA PVT. LTD.
2ND FLOOR 65-A, OMPRO TOWER, KALU SARAI, NEW DELHI, DELHI 110016

Geometric Representation

Class 10 | Mathematics

04 Real Numbers



Misostudy.com



Scientific eBook

This is a sample eBook. The eBook provides chapter-wise theories and examples, formulas, diagrams, and Exam pattern Problem-Solving of the full syllabus.



Complete video lectures

The theory and Problem-Solving in this eBook provide specially designed the video lectures by experienced faculties. Get Free trial and preview on Misostudy.com.

01. Euclid's Division Lemma

Euclid's Division Lemma : Let a and b be any two positive integers. Then, there exist unique integers q and r such that

$$a = bq + r, 0 \leq r < b$$

If $b \mid a$, then $r = 0$. Otherwise, r satisfies the stronger inequality $0 < r < b$.

Remark I The above Lemma is nothing but a restatement of the long division process we have been doing all these years, and that the integers q and r are called the quotient and remainder, respectively.

Remark II The above Lemma has been stated for positive integers only. But, it can be extended to all integers as stated below :

Let a and b any two integers with $b \neq 0$. Then, there exist unique integers q and r such that

$$a = bq + r, \text{ where } 0 \leq r < |b|$$

Remark III

- (i) When a positive integer is divided by 2, the remainder is either 0 or 1. So, any positive integer is of the form $2m$, $2m + 1$ for some integer m .
- (ii) When any positive integer is divided by 3, the remainder is 0 or 1 or 2. So, any positive integer can be written in the form $3m$, $3m + 1$ or $3m + 2$ for some integer m .
- (iii) When a positive integer is divided by 4, the remainder can be 0 or 1 or 2 or 3. So, any positive integer is of the form $4q$ or, $4q + 1$ or, $4q + 2$ or, $4q + 3$.

Example Show that any positive odd integer is of the form $4q + 1$ or $4q + 3$, where q is some integer.

Solution Let a be any odd positive integer and $b = 4$. By division Lemma there exists integers q and r such that

$$a = 4q + r, \text{ where } 0 \leq r < 4$$

$$\Rightarrow a = 4q \text{ or, } a = 4q + 1 \text{ or, } a = 4q + 2 \text{ or, } a = 4q + 3$$

$$[\because 0 \leq r < 4 \Rightarrow r = 0, 1, 2, 3]$$

$$\Rightarrow a = 4q + 1 \text{ or, } a = 4q + 3$$

$$[\because a \text{ is an odd integer } \therefore a \neq 4q, a \neq 4q + 2]$$

Hence, any odd integer is of the form $4q + 1$ or, $4q + 3$.

02. Euclid's Division Algorithm

In order to compute the HCF of two positive integers, say a and b , with $a > b$ we may follow the following steps :

Step I Apply Euclid's division lemma to a and b and obtain whole numbers q_1 and r_1 such that $a = bq_1 + r_1$, $0 \leq r_1 < b$.

Step II If $r_1 = 0$, b is the HCF of a and b

Step III If $r_1 \neq 0$, apply Euclid's division lemma to b and r_1 and obtain two whole numbers q_2 and r_2 such that $b = r_1q_2 + r_2$.

Step IV If $r_2 = 0$, then r_2 is the HCF of a and b .

Step V If $r_2 \neq 0$, then apply Euclid's division lemma to r_1 and r_2 and continue the above

process till the remainder r_n is zero. The divisor at this stage i.e. r_{n-1} , or the non-zero remainder at the previous stage, is the HCF of a and b .

Example Use Euclid's division algorithm to find the HCF of 4052 and 12576.

Solution Given integers are 4052 and 12576 such that $12576 > 4052$. Applying Euclid's division lemma to 12576 and 4052, we get

$$12576 = 4052 \times 3 + 420 \quad \dots(i) \quad \left[\begin{array}{r} 3 \\ \therefore 4052 \overline{)12576} \\ \underline{12156} \\ 420 \end{array} \right]$$

Since the remainder $420 \neq 0$. So, we apply the division lemma to 4052 and 420, to get

$$4052 = 420 \times 9 + 272 \quad \dots(ii) \quad \left[\begin{array}{r} 9 \\ \therefore 420 \overline{)4052} \\ \underline{3780} \\ 272 \end{array} \right]$$

We consider the new divisor 420 and the new remainder 272 and apply division lemma to get

$$420 = 272 \times 1 + 148 \quad \dots(iii) \quad \left[\begin{array}{r} 1 \\ \therefore 272 \overline{)420} \\ \underline{272} \\ 148 \end{array} \right]$$

Let us now consider the new divisor 272 and the new remainder 148 and apply division lemma to get

$$272 = 148 \times 1 + 124 \quad \dots(iv) \quad \left[\begin{array}{r} 1 \\ \therefore 148 \overline{)272} \\ \underline{148} \\ 124 \end{array} \right]$$

We consider now the new divisor 148 and the new remainder 124 and apply division lemma to get

$$148 = 124 \times 1 + 24 \quad \dots(v) \quad \left[\begin{array}{r} 1 \\ \therefore 124 \overline{)148} \\ \underline{124} \\ 24 \end{array} \right]$$

We consider now the new divisor 124 and the new remainder 24 and apply division lemma to get

We consider the new divisor 24 and the new remainder 4 and apply division lemma to get

$$24 = 4 \times 6 + 0 \quad \dots(\text{vii}) \quad \left[\begin{array}{r} 6 \\ \cdot\cdot 4 \overline{) 24} \\ \underline{24} \\ 0 \end{array} \right]$$

We observed that the remainder at this stage is zero. Therefore, the divisor at this stage i.e. 4 (or the remainder at the earlier stage) is the HCF of 4052 and 12576.

Example Any contingent of 616 members is to march behind an army band of 32 members in a parade. The two groups are to march in the same number of columns. What is the maximum number of columns in which they can march?

Solution The maximum number of columns is the HCF of 616 and 32. In order to find the HCF of 616 and 32, let us apply Euclid's division lemma to 616 and 32 to get

$$616 = 32 \times 19 + 8$$

Let us now take the divisor 32 as dividend and remainder 8 as divisor and apply Euclid's division lemma to get

$$32 = 8 \times 4 + 0$$

Since, the remainder at this stage is 0. Therefore, the last divisor i.e. 8 is the HCF of 616 and 32.

Hence, the maximum number of columns in which they can march is 8.

03. The Fundamental Theorem Arithmetic

Result Fundamental Theorem of Arithmetic : Every composite number can be expressed (factorised) as a product of primes, and this factorization is unique except for the order in which the prime factors occur.

Result Let p be a prime number and a be a positive integer. If p divides a^2 , then p divides a .

Example Prove that there is no natural number for which 4^n ends with the digit zero.

Solution We know that any positive integer ending with the digit zero is divisible by 5 and so its prime factorization must contain the prime 5.

$$4^n = (2^2)^n = 2^{2n}$$

\Rightarrow The only prime in the factorization of 4^n is 2.

\Rightarrow There is no other primes in the factorization of $4^n = 2^{2n}$.

[By uniqueness of the Fundamental Theorem of Arithmetic]

\Rightarrow 5 does not occur in the prime factorization of 4^n for any n .

\Rightarrow 4^n does not end with the digit zero for any natural number n .

04. Some Applications of the Fundamental Theorem of Arithmetic

(i) Finding HCF and LCM of Positive Integers

In order to find the HCF and LCM of two or more positive integers, we may use the following algorithm.

Algorithm

Step I Factorize each of the given positive integers and express them as a product of powers of primes in ascending order of magnitudes of primes.

Step II To find the HCF, identify common prime factors and find the smallest (least) exponent of these common factor, Now raise these common prime factors to their smallest exponents and multiply them to get the HCF.

To find the LCM, list all prime factors (once only) occurring in the prime factorisation of the given positive integers.

For each of these factors, find the greatest exponent and raise each prime factor to the greatest exponent and multiply them to get the LCM.

Remark To find the LCM of two positive integers a and b , we can also use the following result, if we have already found the HCF.

$$HCF \times LCM = a \times b.$$

Example Find the HCF and LCM of 144, 180 and 192 by prime factorisation method.

Solution Using the factor tree for the prime factorisation of 144, 180 and 192, we have

$$144 = 2^4 \times 3^2, 180 = 2^2 \times 3^2 \times 5 \text{ and } 192 = 2^6 \times 3$$

To find the HCF, we list the common prime factors and their smallest exponents in 144, 180 and 192 as follows :

Common prime factors	Least exponents
2	2
3	1

$$\therefore HCF = 2^2 \times 3^1 = 12$$

To find the LCM, we list all prime factors of 144, 180, 192 and their greatest exponents as follows :

Prime factors of 144, 180 and 192	Greatest exponents
2	6
3	2
5	1

$$\therefore LCM = 2^6 \times 3^2 \times 5^1 = 64 \times 9 \times 5 = 2880$$

(ii) Proving Irrationality of Numbers

Example Prove that $\sqrt{3}$ is an irrational number.

Solution Let us assume on the contrary that $\sqrt{3}$ is a rational number. Then, there exist positive integers a and b such that

$$\sqrt{3} = \frac{a}{b}, \text{ where } a \text{ and } b \text{ are co-prime i.e. their HCF is 1.}$$

Solution Now,

$$\begin{aligned} \sqrt{3} &= \frac{a}{b} \\ \Rightarrow 3 &= \frac{a^2}{b^2} \\ \Rightarrow 3b^2 &= a^2 \\ \Rightarrow 3 &| a^2 && [\because 3 | 3b^2] \\ \Rightarrow 3 &| a && \dots(i) \\ \Rightarrow a &= 3c \text{ for some integer } c \\ \Rightarrow a^2 &= 9c^2 \\ \Rightarrow 3b^2 &= 9c^2 && [\because a^2 = 3b^2] \\ \Rightarrow b^2 &= 3c^2 \\ \Rightarrow 3 &| b^2 && [\because 3 | 3c^2] \\ \Rightarrow 3 &| b && \dots(ii) \end{aligned}$$

From (i) and (ii), we observe that a and b have at least 3 as a common factor. But this contradicts the fact that a and b are co-prime. This means that our assumption is not correct.

Hence, $\sqrt{3}$ is an irrational number.

Example Prove that $5 - \sqrt{3}$ is an irrational number.

Solution Let us assume on the contrary that $5 - \sqrt{3}$ is rational. Then, there exist co-prime positive integers a and b such that

$$\begin{aligned} 5 - \sqrt{3} &= \frac{a}{b} \\ \Rightarrow 5 - \frac{a}{b} &= \sqrt{3} \\ \Rightarrow \frac{5b - a}{b} &= \sqrt{3} \\ \Rightarrow \sqrt{3} &\text{ is rational } [\because a, b \text{ are integers } \therefore \frac{5b - a}{b} \text{ is a rational number}] \end{aligned}$$

This contradicts the fact $\sqrt{3}$ is irrational. So, our assumption is incorrect.

Hence, $5 - \sqrt{3}$ is an irrational number.

(iii) Determining the Nature of the Decimal Expansions of Rational Numbers

Result Let x be a rational number whose decimal expansion terminates. Then, x can

expressed in the form $\frac{p}{q}$, where p and q are co-primes, and the prime factorisation of q is of the form $2^m \times 5^n$, where m, n are non-negative integers.

Result Let $x = \frac{p}{q}$ be a rational number, such that the prime factorisation of q is of the form $2^m \times 5^n$, where m, n are non-negative integers. Then, x has a decimal expansion which terminates after k places of decimals, where k is the larger of m and n .

CBSE Pattern Exercise (1)

(Q 1 to 2) One Mark

- Use Euclid's division algorithm to find the HCF of : 196 and 38220
- Find the LCM and HCF of the following pair of integers and verify that $\text{LCM} \times \text{HCF} =$ product of the two numbers. 510 and 92.

(Q 3 to 6) Two Marks

- Given that $\text{HCF}(306, 657) = 9$, find $\text{LCM}(306, 657)$.
- Prove that the following are irrational : $6 + \sqrt{2}$
- Without actually performing the long division, state whether the following rational number will have a terminating decimal expansion or a non-terminating repeating decimal expansion: $\frac{23}{2^3 5^2}$.
- Without actually performing the long division, state whether the following rational number will have a terminating decimal expansion or a non-terminating repeating decimal expansion: $\frac{77}{210}$

(Q 7 to 10) Four Marks

- Use Euclid's division lemma to show that the cube of any positive integer is of the form $9m$, $9m + 1$ or $9m + 8$.
- Prove that $\sqrt{5}$ is irrational.
- The following real number have decimal expansions as given below. Decide whether it is rational or not. If it is rational, and of the form $\frac{p}{q}$, what can you say about the prime factors of q ? 43.123456789.
- The following real number have decimal expansions as given below. Decide whether it is rational or not. If it is rational, and of the form $\frac{p}{q}$, what can you say about the prime factors of q ? 0.120120012000120000...



Answer & Solution

Q1

Since $38220 > 196$, we apply the division lemma to 38220 and 196 to obtain

$$38220 = 196 \times 195 + 0.$$

Since the remainder is zero, the process stops. Since the divisor at this stage is 196, Therefore, HCF of 196 and 38220 is 196.

Q2

510 and 92

$$510 = 2 \times 3 \times 5 \times 17$$

$$92 = 2 \times 2 \times 23$$

$$\text{HCF} = 2$$

$$\text{LCM} = 2 \times 2 \times 3 \times 5 \times 17 \times 23 = 23460$$

$$\text{Product of the two numbers} = 510 \times 92 = 46920$$

$$\begin{aligned} \text{HCF} \times \text{LCM} &= 2 \times 23460 \\ &= 46920 \end{aligned}$$

Hence, product of two numbers = HCF \times LCM

Q3

We know that, $\text{LCM} \times \text{HCF} = \text{Product of two numbers}$

$$\therefore \text{LCM} \times \text{HCF} = 306 \times 657$$

$$\text{LCM} = \frac{306 \times 657}{\text{HCF}} = \frac{306 \times 657}{9}$$

$$\text{LCM} = 22338$$

Q4

Let $6 + \sqrt{2}$ be rational.

Therefore, we can find two integers a, b ($b \neq 0$) such that

$$6 + \sqrt{2} = \frac{a}{b}$$

$$\sqrt{2} = \frac{a}{b} - 6$$

since a and b are integers, $\frac{a}{b} - 6$ is also rational and hence, $\sqrt{2}$ should be rational. This

04 Real Numbers

contradicts the fact that $\sqrt{2}$ is irrational. Therefore, our assumption is false and hence, $6 + \sqrt{2}$ is irrational.

Q5

$$\frac{23}{2^3 \times 5^2}$$

The denominator is of the form $2^m \times 5^n$.

Hence, the decimal expansion of $\frac{23}{2^3 \times 5^2}$ is terminating.

Q6

$$\frac{77}{210} = \frac{11 \times 7}{30 \times 7} = \frac{11}{30}$$

$$30 = 2 \times 3 \times 5$$

since the denominator is not of the form $2^m \times 5^n$, and it also has 3 as its factors, the decimal expansion of $\frac{77}{210}$ is non-terminating repeating.

Q7

Let a be any positive integer and $b = 3$

$a = 3q + r$, where $q \geq 0$ and $0 \leq r < 3$

$\therefore a = 3q$ or $3q + 1$ or $3q + 2$

Therefore, every number can be represented as these three forms.

There are three cases.

Case I When $a = 3q$,

$$a^3 = (3q)^3 = 27q^3 = 9(3q^3) = 9m,$$

Where m is an integer such that $m = 3q^3$

Case II When $a = 3q + 1$

$$a^3 = (3q + 1)^3$$

$$a^3 = 27q^3 + 27q^2 + 9q + 1$$

$$a^3 = 9(3q^3 + 3q^2 + q) + 1$$

$$a^3 = 9m + 1$$

Where m is an integer such that $m = (3q^3 + 3q^2 + q)$

Case III When $a = 3q + 2$,

$$a^3 = (3q + 2)^3$$

$$a^3 = 27q^3 + 54q^2 + 36q + 8$$

$$a^3 = 9(3q^3 + 6q^2 + 4q) + 8$$

$$a^3 = 9m + 8$$

Where m is an integer such that $m = (3q^3 + 6q^2 + 4q)$

Therefore, the cube of any positive integer is of the form $9m$, $9m + 1$, or $9m + 8$.

Q8

Let $\sqrt{5}$ is a rational number.

Therefore, we can find two integers $a, b (b \neq 0)$ such that $\sqrt{5} = \frac{a}{b}$

Let a and b have a common factor other than 1. Then we can divide them by the common factor, and assume that a and b are co-prime.

$$a = \sqrt{5}$$
$$a^2 = 5b^2$$

Therefore, a^2 is divisible by 5 and it can be said that a is divisible by 5.

Let $a = 5k$, where k is an integer

$$(5k)^2 = 5b^2$$

$b^2 = 5k^2$ This means that b^2 is divisible by 5 and hence, b is divisible by 5.

This implies that a and b have 5 as a common factor.

And this is a contradiction to the fact that a and b are co-prime.

Hence, $\sqrt{5}$ cannot be expressed as $\frac{p}{q}$ or it can be said that $\sqrt{5}$ is irrational.

Q9

Since this number has a terminating decimal expansion, it is a rational number of the form $\frac{p}{q}$ and q is of the form $2^n \times 5^n$ i.e., the prime factors of q will be either 2 or 5 or both.

Q10

The decimal expansion is neither terminating nor recurring. Therefore, the given number is an irrational number.

miso
study

CBSE Exam Pattern Exercise

Objective Questions (1)

- the decimal expansion of number $\frac{441}{2^2 \times 5^3 \times 7}$ has
 - a terminating decimal
 - non-terminating but repeating
 - non-terminating non-repeating
 - terminating after two decimal places
- n^2-1 is divisible by 8, if n is
 - an integer
 - a natural number
 - an odd integer
 - an even integer
- The HCF of 95 and 152 is
 - 57
 - 1
 - 19
 - 38
- If two positive integers a and b are written as $a = x^3y^2$ and $b = xy^3$; x, y are prime numbers, then HCF = (a, b) is _____.
 - xy
 - xy^2
 - x^3y^3
 - x^2y^2
- The decimal expansion of the rational number $\frac{33}{2^2 \times 5}$ will terminate after
 - one decimal place
 - two decimal places
 - three decimal places
 - more than 3 decimal places
- If the sum of LCM and HCF of two numbers is 1260 and their LCM is 900 more than their HCF, then the product of two numbers is _____.
 - 203400
 - 194400
 - 198400
 - 205400

7. If $a = 2^3 \times 3$, $b = 2 \times 3 \times 5$, $c = 3^n \times 5$ and $\text{LCM}(a, b, c) = 2^3 \times 3^2 \times 5$, then $n =$ _____.
- (a) 1
(b) 2
(c) 3
(d) 4
8. The largest number which divides 70 and 125, leaving remainders 5 and 8, respectively is
- (a) 13
(b) 65
(c) 875
(d) 1750
9. If the LCM of a and 18 is 36 and the HCF of a and 18 is 2, then $a =$ _____.
- (a) 2
(b) 3
(c) 4
(d) 1
10. If the HCF of 65 and 117 is expressible in the form $65m - 117$, then the value of m is _____.
- (a) 4
(b) 2
(c) 1
(d) 3



Answer & Solutions

1. (a)

$$\frac{441}{2^3 \times 5^3 \times 7} \quad (\text{on simplification})$$

$$= \frac{63}{2^3 \times 5^3}$$

Since the denominator only contains the power of 2 and 5

$\therefore \frac{441}{2^3 \times 5^3 \times 7}$, on simplification gives a terminating decimal expansion

2. (c)

$$\text{Let } a = n^2 - 1$$

Here n can be even or odd

case1:- n = even ie n = 2k where k is any integer

$$a = (2k)^2 - 1$$

$$a = 4k^2 - 1$$

when R = -1, $4(-1)^2 - 1 = 3$, not divisible by 8.

when R = 0, $4(0)^2 - 1 = -1$, not divisible by 8.

case2:- n = odd, n = 2k + 1, where k is any integer

$$a = (2R + 1)^2 - 1$$

$$a = 4k^2 + 1 + 4k - 2 \quad [\text{using } (a + b)^2]$$

$$a = 4k^2 + 4k$$

$$\text{When } R = -1; \quad a = 3(-1)^2 + 4(-1)$$

$$= 4 - 4 = 0 \text{ which is divisible by 8}$$

$$\text{When, } R = 0; \quad a = 4(0)^2 + 4(0) = 0 \text{ which is divisible by 8.}$$

$$\text{When, } R = 1; \quad a = 4(1)^2 + 4(1) = 8 \text{ which is divisible by 8.}$$

\Rightarrow n is an odd number.

3. (c)

HCF (95,152)

By using Euclid's division lemma

$$a = bq + r; \quad 0 \leq r < b$$

where a = dividend; b = divisor; q = quotient; r = remainder

$$152 = 95 \times 1 + 57$$

$$95 = 57 \times 1 + 38$$

$$57 = 38 \times 1 + 19$$

$$38 = 19 \times 2 + 0$$

$$\text{HCF (a,b)} = 19$$

04 Real Numbers

4. (b)

$$a = x^3 y^2 ; b = xy^3$$

HCF = The lowest of indices of x and y

$$\text{HCF}(a,b) = x y^2$$

5. (b)

$$\frac{33}{2^2 \times 5}$$

on simplifying the denominator

$$\frac{33}{2^2 \times 5} = \frac{33}{4 \times 5} = \frac{33}{20} = 1.65$$

\therefore It terminates after 2 decimal places.

6. (b)

According to the question,

$$\text{LCM} + \text{HCF} = 1260 \quad (1)$$

also, LCM is 900 more than their HCF

$$\Rightarrow \text{HCF} + 900 = \text{LCM} \quad (2)$$

Substitute (2) in (1)

$$\text{HCF} + 900 + \text{HCF} = 1260$$

$$2 \text{ HCF} = 1260 - 900$$

$$\text{HCF} = 180$$

$$\text{Put HCF} = 180 \text{ in (2)}$$

$$\text{LCM} = 180 + 900 = 1080$$

Since the product of LCM and HCF of true numbers is equal to the product of numbers.

$$\Rightarrow \text{Product of numbers} = 180 \times 1080 = \underline{194400}$$

7. (b)

$$a = 2^3 \times 3 ; b = 2 \times 3 \times 5 ; c = 3^n \times 5$$

$$\text{LCM}(a,b,c) = 2^3 \times 3^2 \times 5$$

Also, LCM = Highest indices of 2,3 and 5.

$$\Rightarrow n = 2$$

8. (a)

To find the largest numbers which divides 70 and 125 leaving the remainder 5 and 8 respectively, We will subtract 5 from 70 and 8 from 125 then we will compute the HCF of the result obtained after subtraction.

$$70 - 5 = 65 ; 125 - 8 = 117$$

$$65 = 13 \times 5$$

$$117 = 13 \times 3^2$$

$$\text{HCF}(65, 117) = 13$$

\therefore 13 is the largest number which divides 70 & 125 leaving the remainder 5 and 8 respectively

9. (c)

Product of number = LMC \times HCF of the numbers.

$$18 \times a = 36 \times 2$$

$$a = \frac{236 \times 2}{18}$$

$$a = 4$$

10.(b)

HCF (65,117)

$$65 = 13 \times 5 \quad ; \quad 117 = 13 \times 3^2$$

HCF(65, 117) = 13

65 = 117 = 13 as HCF (65, 117) is expressible as $65m - 117$

$$65m = 13 + 117$$

$$65m = 130$$

$$m = \frac{130}{65} = 2 \Rightarrow m = 2$$



misostudy